

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A01N 63/00, A61K 35/14, 38/00, C07H 17/00, C07K 14/00, C12N 1/00, 5/00, 5/06, 5/22, 7/01, 15/00, 15/03, 15/09, 15/06, 15/11, C12P 1 /00, 21 /06	A1	(11) International Publication Number: WO 95/23512 (43) International Publication Date: 8 September 1995 (08.09.95)
(21) International Application Number: PCT/US95/02944 (22) International Filing Date: 1 March 1995 (01.03.95) (30) Priority Data: 08/205,720 3 March 1994 (03.03.94) US (71) Applicant: ALEXION PHARMACEUTICALS, INC. [US/US]; Suite 360, 25 Science Park, New Haven, CT 06511 (US). (72) Inventors: ROTHER, Russell, P.; 384 Black's Road, Cheshire, CT 06410 (US). ROLLINS, Scott; 12 Nutmeg Circle, Monroe, CT 06468 (US). SQUINTO, Stephen, P.; 16 Coachmans Lane, Bethany, CT 06524 (US). (74) Agent: KLEE, Maurice, M.; 1951 Burr Street, Fairfield, CT 06430 (US).		(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: TERMINAL COMPLEMENT INHIBITOR FUSION GENES AND PROTEINS (57) Abstract Nucleic acid sequences encoding chimeric proteins that comprise a functional portion of a parent terminal complement inhibitor, such as CD59, and a heterologous transmembrane domain are provided. The parent terminal complement inhibitor is modified to inactivate its GPI signal sequence. The heterologous transmembrane domain serves to anchor the chimeric protein to the cell membrane without substantially interfering with the complement inhibitor activity of the terminal complement inhibitor. The nucleic acid sequences and encoded chimeric proteins can be used to protect cells from complement attack.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

5 TERMINAL COMPLEMENT INHIBITOR FUSION GENES AND PROTEINSFIELD OF THE INVENTION

 The present invention relates to terminal complement
inhibitory proteins that have been genetically engineered
10 to alter their attachment to the cell surface and to
medical uses of such novel molecules.

BACKGROUND OF THE INVENTIONI. The Complement System

 The complement system acts in conjunction with other
15 immunological systems of the body to defend against
intrusion of cellular and viral pathogens. There are at
least 25 complement proteins, which are found as a
complex collection of plasma proteins and membrane
cofactors. The plasma proteins make up about 10% of the
20 globulins in vertebrate serum. Complement components
achieve their immune defensive functions by interacting
in a series of intricate but precise enzymatic cleavage
and membrane binding events. The resulting complement
cascade leads to the production of products with opsonic,
25 immunoregulatory, and lytic functions.

 The lytic aspect of complement function is effected
by the permeabilization of target cell membranes as a
direct action of an assemblage of complement proteins
known individually as "terminal complement components"
30 and, in their functional assemblage, as the membrane
attack complex, or "MAC". (See Esser, 1991; and Bhakdi,
et al., 1991.) The actions of the MAC, hereinafter
referred to as "complement attack," create pores or leaky
patches that lead to the disruption of osmotic and ionic
35 gradients in target cells, which, at high enough MAC
concentrations, causes cell death. Lower concentrations
of MACs can produce other effects, including activation

-2-

of endothelial cells and platelets. Inappropriate MAC activity can result in pathologic damage to cells and tissues.

5 The complement cascade progresses via the classical pathway or the alternative pathway. These pathways share many components, and, while they differ in their early steps, both converge and share the same terminal complement components responsible for complement attack and the activation and/or destruction of target cells.

10 The classical complement pathway is typically initiated by antibody recognition of and binding to an antigenic site on a target cell. The alternative pathway is usually antibody independent, and can be initiated by certain molecules on pathogen surfaces. Both pathways
15 converge at the point where complement component C3 is cleaved by an active protease (which is different in each pathway) to yield C3a and C3b. Other pathways activating complement attack can act later in the sequence of events leading to various aspects of complement function,
20 including the formation of the MAC.

C3a is an anaphylatoxin that can induce degranulation of mast cells, resulting in the release of histamine and other mediators of inflammation. C3b has multiple functions. As opsonin, it binds to bacteria,
25 viruses and other cells and particles and tags them for removal from the circulation. C3b can also form a complex with other components unique to each pathway to form classical or alternative C5 convertase, which cleaves C5 into C5a (another anaphylatoxin), and C5b,
30 which is the first of the terminal complement components that make up the MAC. (Amongst the several means by which complement attack can be initiated, proteolytic enzymes with relatively broad target protein specificities, including plasmin, elastase, and cathepsin G, can cleave
35 C5 so as to mimic the action of C5 convertase and produce active C5b.) C5b combines sequentially with C6, C7, and C8 to form the C5b-8 complex at the surface of the target

-3-

cell. Upon binding of several C9 molecules, the active MAC (C5b-9) is formed.

II. Regulation of the Complement System

5 Normally, the complement system is in a continuous state of spontaneous turnover. C3 can spontaneously acquire C3b functions, forming a functional C3 convertase and leading to the formation of more C3b. The C3b generated in this spontaneous fashion can also form C5 convertase and thus initiate the final steps in the cascade that forms the MAC.

10 Under normal conditions, blood flow will dilute and disperse the low levels of spontaneously activated complement components thus helping to prevent MAC buildup in any one location in the vasculature. In addition, homeostatic regulation of the actions of autologous complement proteins to prevent autoimmune attack is mediated by specific endogenous complement inhibitor proteins (CIPs), that can be found on the surfaces of most human cells. Ordinarily, blood flow and the action of CIPs suffice to render cells resistant to normal levels of spontaneous complement activation without injury or lysis. Under conditions of acute inflammation, and in various disease states where complement activation and MAC formation are accelerated, the normal quantity and activity of endogenous complement inhibitors may be inadequate to protect autologous cells from MAC-induced lysis and/or sublytic MAC-induced cell activation. Endogenous CIP activity may also be insufficient where there is stasis of the blood, and/or where there are defects in or deficiencies of naturally occurring inhibitors.

25 A number of CIPs have been identified that serve to protect cells from damage mediated by complement from concordant species. See Zalman, et al., 1986; Schonermark, et al., 1986; Nose, et al., 1990; and Sugita, et al., 1988. These inhibitors act at various defined points in the complement cascade. For example,

CD55, also known as decay accelerating factor (DAF), exerts its major inhibitory effects on the actions of C3 convertase.

5 In cases where the complement cascade is initiated at points in the pathway after the C3 convertase step, such as through the generation of active C5b by broad spectrum proteases, DAF and other complement inhibitors acting at earlier steps in the cascade sequence are ineffective. There are, however, inhibitors that do not
10 share this deficiency. These inhibitors act at the final steps in MAC assembly and thus can effectively block complement attack initiated by almost any means. These inhibitors are known as "terminal complement inhibitors" or "terminal CIPs."

15 III. Terminal CIPs

The most thoroughly characterized terminal CIP is the human protein CD59 (also known as "protectin", "MACIF", or "p18"). CD59 is a glycoprotein with an apparent molecular mass of 18-21 kilodaltons that
20 protects cells from complement-mediated lysis. CD59 is tethered to the outside of the cell by a glycosyl-phosphatidylinositol (GPI) glycolipid moiety that anchors it in the cell membrane. CD59 is found associated with the membranes forming the surfaces of
25 most human cells including erythrocytes, lymphocytes, and vascular endothelial cells. (See, for example, Sims, et al., U.S. Patent No, 5,135,916.)

CD59 appears to function by competing with C9 for binding to C8 in the C5b-8 complex, thereby decreasing
30 the formation of the C5b-9 MAC (Rollins, et al., 1990). CD59 thus acts to reduce both cell stimulation and cell lysis by MACs (Rollins, et al., 1990; Rollins, et al., 1991; Stefanova, et al., 1989; Sugita, et al., 1988; Davies, et al., 1989; Holguin, et al., 1989; Okada, et
35 al., 1989a; Meri, et al., 1990; Whitlow, et al., 1990; and Harada, et al., 1990). This activity of CD59 is for the most part species-selective, most efficiently

blocking the formation of MACs under conditions where C8 and C9 are derived from homologous (i.e., human) serum (Venneker, et al., 1992). The assimilation of purified CD59 into the plasma membrane of non-human erythrocytes (which are believed to be protected from homologous non-human complement attack by the action of their own cell surface complement inhibitor proteins) and oligodendrocytes (brain cells which are believed to be protected less, if at all, by cell surface proteins, but may be protected in vivo by the blood brain barrier) has shown that CD59 can protect these cells from cell lysis mediated by human complement. (Rollins, et al., 1990; Rollins, et al., 1991; Stefanova, et al., 1989; Meri, et al., 1990; Whitlow, et al., 1990; Okada, et al., 1989b; and Wing, et al., 1992).

cDNAs coding for CD59 have been cloned and the structure of the CD59 gene has been characterized (Davies, et al., 1989; Okada, et al., 1989b; Philbrick, et al., 1990; Sawada, et al., 1989; and Tone, et al., 1992). Non-human mammalian cells transfected with the cloned CD59 cDNA, and thereby expressing the human CD59 protein on their cell surfaces, have been shown to gain resistance to complement-mediated cell lysis (Zhao, et al., 1991; and Walsh, et al., 1991).

CD59 has been reported to be structurally related to the murine Ly-6 antigens (Philbrick, et al., 1990; and Petranksa, et al., 1992). The genes encoding these antigens, also known as T-cell activating proteins, are members of the Ly-6 multigene family, and include Ly-6A.2, Ly-6B.2, Ly-6C.1, Ly6C.2, and Ly-6E.1. The gene encoding the murine thymocyte B cell antigen ThB is also a member of this family (Shevach, et al. 1989; and Gumley, et al., 1992).

A number of viral and non-human primate complement inhibitor proteins that are similar in structure and function to CD59 have been described (see Rother, et al., 1994; Albrecht, et al., 1992; commonly assigned,

compending, U.S. patent application Serial No. 08/105,735, filed August 11, 1993, by William L. Fodor, Scott Rollins, Russell Rother, and Stephen P. Squinto, and entitled "Complement Inhibitor Proteins of Non-Human Primates;" and commonly assigned and compending PCT patent application Serial No. PCT/US93/00672, filed January 12, 1993, by Bernhard Fleckenstein and Jens-Christian Albrecht, and entitled "Complement Regulatory Proteins of Herpesvirus Saimiri".

5 These proteins -- BABCIP (SEQ ID NO:1), AGMCIP (SEQ ID NO:2), SQMCIP (SEQ ID NO:3), OWMCIP (SEQ ID NO:4), MARCIP (SEQ ID NO:5), and HVS-15 (SEQ ID NO:6) -- all share striking sequence homologies, including a distinctive conserved arrangement of cysteines within
10 their amino acid sequences. These conserved patterns are most readily perceived by aligning the sequences of the proteins so that the cysteine residues are in register as seen in FIG. 1.

15 Cysteine residues of many proteins form a structural element referred to in the art as a "cysteine backbone". In proteins in which they occur, cysteine backbones play essential roles in determining the three-dimensional folding, tertiary structure, and ultimate function of the molecule. The proteins of the Ly-6 multigene family, as
20 well as several other proteins, share a particular cysteine backbone structure referred to herein as the "Ly- 6 motif". For example, the human urokinase plasminogen activator receptor (uPAR; Roldan, et al., 1990) and one of several squid glycoproteins of unknown
25 function (Sgp2; Williams, et al., 1988) contain the Ly-6 motif.


30 Subsets of proteins having the Ly-6 motif can be identified by the presence of conserved amino acid residues immediately adjacent to the cysteine residues.
35 Such conservation of specific amino acids within a subset of proteins can be associated with specific aspects of the folding, tertiary structure, and ultimate function of

-7-


the proteins. These conserved patterns are most readily perceived by aligning the sequences of the proteins so that the cysteine residues are in register.

As discussed fully in the above-referenced,
5 copending U.S. patent application Serial No. 08/105,735, the relevant portions of which are incorporated herein by reference, a series of non-human primate C5b-9 inhibitory proteins have been identified which are characterized by a cysteine backbone structure which defines a specific
10 subset of the general Ly-6 motif.

Specifically, these non-human primate CIPs include polypeptides comprising a cysteine backbone with a Ly-6 motif characterized by the formula:



In addition, the non-human primate C5b-9 inhibitory proteins include amino acid sequences conforming to the following formula:



In both formulas, the X in X_n indicates a peptide containing any combination of amino acids, the n in X_n represents the length in amino acid residues of the
25 peptide, and each X at any position can be the same as or different from any other X of the same length in any other position.

As discussed fully in the above-referenced, copending, commonly assigned, PCT application Serial No.
30 PCT/US 93/00672, the relevant portions of which are incorporated herein by reference, and in Albrecht, et al., 1992, a protein of the herpesvirus saimiri having C5b-9 inhibitory activity has been discovered (referred to herein as "HVS-15"). This viral protein has the Ly-6
35 motif which is characteristic of the non-human primate C5b-9 inhibitory proteins discussed above, i.e., its structure is described by formulas (1) and (2) above.

In the discussion which follows, terminal CIPs comprising Ly-6 motifs are referred to as "Ly-6 terminal CIPs." These CIPs will in general satisfy formula (1) above and preferably also formula (2). Some variations, however, in the spacing between any two of the ten cysteines making up the Ly-6 motif and in the adjacent amino acids are to be expected in as yet uncharacterized terminal CIPs of other species.

Also, Petranka et al., 1993, and Norris, et al., 1993, have reported that in CD59 (SEQ ID NO:7) the disulfide bond between Cys6 and Cys13, as well as the disulfide bond between Cys64 and Cys69, can be disrupted by replacement of these cysteines with serines without substantially compromising the functionality of CD59. These cysteines correspond to the second, third, ninth, and tenth cysteines in the above formulas. Accordingly, as used herein, the term "Ly-6 terminal CIP" is intended to also include terminal complement inhibitor proteins conforming with the above formulas but with all or some of the second, third, ninth or tenth cysteines replaced with serine, or another amino acid.

IV. Other Cell Surface Complement Inhibitor Proteins

In addition to the Ly-6 terminal CIPs discussed above, other membrane bound CIPs have been described in the literature, including the following:

(a) CD46 (membrane cofactor protein, MCP, see, for example, PCT patent publication No. WO 91/02002) is a 350 amino acid transmembrane (TM) protein found on all cells except red blood cells. CD46 binds to C3b, and, once bound, promotes the activity of proteases that cleave C3b into inactive fragments, thus preventing C3b accumulation on the cell surface and, in turn, protecting cells from complement attack. Both membrane bound and secreted forms of CD46 have been reported in the literature (Purcell et al., 1991).

(b) CD55 (decay accelerating factor, DAF), mentioned above, is a GPI-anchored cell surface protein

present on all cells including red blood cells. Unlike CD46, CD55 does not destroy C3b. Rather, CD55 prevents C3b from reacting with other complement components, thus contravening complement mediated cytolysis. Both membrane bound and secreted forms of CD55 have been reported in the literature (Moran et al., 1992).

(c) CD35 (complement receptor 1, CR1) is found on a select group of lymphocytes as well as erythrocytes, neutrophils, and eosinophils, and causes degradation of C3b molecules adhering to neighboring cells.

(d) Factor H and C4b-binding protein, both of which inhibit alternative C3 convertase activity.

V. Transplantation

Intrinsic activation of complement attack via the alternative pathway during storage of donor organs is responsible for certain problems associated with organ transplantation which arise as a result of endothelial cell stimulation and/or lysis by the C5b-9 MAC (Brasile, et al. 1985). Ex vivo complement attack leads to reduced vascular viability and reduced vascular integrity when stored organs are transplanted, increasing the likelihood of transplant rejection.

Ten percent of allogeneic transplanted kidneys with HLA-identical matches are rejected by in vivo immunologic mechanisms (Brasile, et al. 1987). In 78% of the patients who reject organs under these conditions, cytotoxic antibodies binding to molecules on the surfaces of vascular endothelial cells are seen (Brasile, et al., 1987). Such antibody cytotoxicity is mediated by complement attack, and is responsible for the rejection of transplanted solid organs including kidneys and hearts (Brasile, et al., 1987; Brasile et al., 1985). Antibody primed, complement-mediated rejection is usually rapid and irreversible, a phenomenon referred to as hyperacute rejection.

In the xenogeneic setting, as when non-human organs are transplanted into human patients, activation of

complement attack by antibodies directed against molecules on the surfaces of endothelial cells lining the vessels of the donor organ is almost always observed. The prevalence of such xenoreactive antibodies accounts for the nearly universal occurrence of hyperacute rejection of xenografts (Dalmasso, et al., 1992). Old world primates, including humans, have high levels of preexisting circulating "natural" antibodies that predominantly recognize carbohydrate determinants expressed on the surface of xenogeneic cells from discordant species. Recent evidence indicates that most of these antibodies react with galactose in an α 1-3 linkage with galactose (Gal(α 1-3)Gal) (Sandrin, et al., 1993).

Old world primates lack the appropriate functional α -1,3-galactose transferase and thus do not express this carbohydrate epitope. Therefore, following transplantation of a vascularized xenogeneic donor organ, these high-titer antibodies bind to the Gal(α 1-3)Gal epitope on the vascular endothelium and activate the recipient's complement through the classical pathway. The massive inflammatory response that ensues from activation of the complement cascade leads to the destruction of the donor organ within minutes to hours.

Xenoreactive antibodies are not exclusively responsible for hyperacute rejection of discordant organs in all cases. For example, erythrocytes from some species can activate human complement via the alternative pathway and newborn piglets raised to be free of preformed antibodies reject xenografts almost immediately. It is therefore likely that in some species combinations, activation of the alternative complement pathway contributes to graft rejection.

Endogenously-expressed, membrane-associated complement inhibitory proteins normally protect endothelial cells from autologous complement. However, the species restriction of complement inhibitors makes them

relatively ineffective with respect to regulating discordant xenogeneic serum complement. The lack of effective therapies aimed at eliminating this antibody and complement-mediated hyperacute rejection presents a major barrier to the successful transplantation of discordant animal organs into human recipients.

Recently, a report on a baboon-to-human liver transplant has been published in which the xenogeneic donor organ failed to exhibit signs of hyperacute rejection (Starzl, et al., 1993). The low levels of anti-baboon antibodies likely to be present in human blood make hyperacute responses less likely. However, it is believed that recently discovered baboon CIPs, which have been shown to be related to CD59 and to be effective against human complement, also played a role in maintaining the integrity of this xenotransplanted organ. (See U.S. patent application Serial No. 08/105,735, referred to above.)

The lack of hyperacute rejection seen in the baboon to human xenotransplant discussed above suggests that complement inhibitor proteins effective against human complement may, in combination with other anti-rejection strategies, allow safe and effective xenotransplantation of transgenic animal organs expressing such proteins into human patients.

VI. GPI-Anchored CIPs and Modifications Thereof

GPI-anchored terminal CIPs share certain properties that make them less desirable than transmembrane (TM) proteins for use as complement inhibiting agents for the protection of transplanted cells or organs.

GPI-anchored terminal CIPs, including CD59, BABCIP, and AGMCIP, can be cleaved from cell surfaces by specific phospholipase enzymes that hydrolyze GPI anchors. Such phospholipases are present in the serum (phospholipase D, Davitz, et al., 1987), and may also be released from cells in response to ischemia (phospholipase C, Vakeva, et al., 1992). Since ischemia is an unavoidable

concomitant of transplantation, the process of transplantation may serve to remove native and/or artificially introduced GPI-anchored terminal CIPs from the very cells within the transplanted organ that they are meant to protect.

Another mechanism by which GPI-anchored proteins are removed from the cell surface is the incorporation of such proteins into membrane vesicles and the subsequent shedding of the vesicles from the cell. Such vesiculation can occur in response to various stimuli, such as ischemia-induced complement attack. It has been reported that GPI-anchored proteins are concentrated in these vesicles relative to their concentration in the cell membrane, a phenomenon that may reflect involvement of these proteins in the vesiculation process itself (Butikofer, et al., 1989; Brown, et al., 1992; Whitlow, et al., 1993). Such preferential incorporation into shed vesicles can reduce the concentrations of GPI-anchored proteins on the cell surface, including the concentrations GPI-anchored terminal CIPs. Such reductions of terminal CIP concentrations, particularly in response to complement attack, may occur at just those times when inhibition of complement is most needed.

In addition to their susceptibility to removal from the cell surface, GPI-anchored proteins also suffer from the problem that their production may be limited in various cell types. That is, only so many GPI-anchored molecules can normally be produced by a cell within a given time frame, so that introducing genes for further GPI-anchored proteins may not in fact result in substantial increases in the amount of protein actually present on the cell surface.

The limiting case of this problem involves cells which are incapable of producing any GPI-anchored proteins. The clinical disease of paroxysmal nocturnal hemoglobinuria (PNH) involves cells of this type, specifically, blood cells which do not produce GPI-

anchored terminal CIPs. As discussed in copending, commonly assigned, U.S. patent application Serial No. 08/206,189, entitled "Method for the Treatment of Paroxysmal Nocturnal Hemoglobinuria", which is being
5 filed concurrently herewith in the names of Russell Rother, Scott A. Rollins, Seth A. Fidel, and Stephen P. Squinto, PNH cells can be made resistant to complement attack through the use of the transmembrane terminal CIPs described herein.

10 A further drawback of GPI-anchored proteins involves the ability of these proteins to transduce signals into the cell upon being cross-linked by specific antibodies and presumably upon binding their natural ligand (Okada, et al., 1989b; Seaman, et al., 1991; Su, et al., 1991;
15 Deckert, et al., 1992; Cinek, et al., 1992; Card, et al., 1991; Groux, et al., 1989; and Stefanova, et al., 1991). Possible undesirable cellular responses to such intracellular signals can include phospholipase activation and/or release, and the stimulation of vesicle
20 formation and shedding, both of which, as discussed above, can result in the loss of GPI-anchored proteins from the cell surface. Thus, the very GPI-anchored terminal CIPs that are used to protect the cells of a transplanted organ from complement attack may activate
25 the cellular events that lead to their removal from the cell surface.

Work has been performed in which the means of attachment of GPI-anchored proteins to the outer cell surface has been varied from their natural GPI anchors by
30 substitution of other anchoring moieties (Su, et al., 1991; and Lublin, et al., 1991).

For example, chimeric derivatives of CD55, containing amino acids 1-304 of CD55 fused to a fragment of CD46 which includes the protein's transmembrane domain
35 (i.e., amino acids 270-350 of CD46) or to a fragment of the human major histocompatibility protein HLA-B44 which includes its transmembrane domain (i.e., amino acids

262-338 of HLA-B44), have been reported to retain levels of function equivalent to native CD55 (Lublin, et al., 1991). Significantly, with regard to the present invention, no such substitutions have been made with terminal CIPs and no such molecules have been developed for clinical use and, in particular, for use in constructing transgenic organs for transplantation.

VII. Protein Structure and Function

Minor alterations of protein primary structures (amino acid sequences) can have profound effects on their functional properties. The best known example of this phenomenon is in the case of sickle cell anemia, in which a single amino acid alteration, namely, a change in residue 6 of the beta chain of hemoglobin from Glu to Val, is sufficient to change the oxygen binding properties of the hemoglobin molecule and to thereby cause sickle cell disease.

The insertion of heterologous amino acid sequences representing new domain structures into a protein can also have significant effects on the protein's functional properties. For example, the introduction of a 10 amino acid epitope of the c-myc proto-oncogene (known as the myc tag) to the int-1 proto-oncogene alters the functional properties of int-1. Specifically, C57MG mammary epithelial cells are transformed by wild-type int-1, but not by the myc-tagged int-1, while residual function of the myc-tagged int-1 gene is seen in a more sensitive assay examining effects on Drosophila development (McMahon et al., 1989).

Additionally, substitution of homologous sequences from heterologous proteins can have profound effects on protein function. For example, replacement of either of the two most carboxyl-terminal 12 amino acid segments of the mouse nerve growth factor gene with homologous segments from the related mouse brain derived neurotrophic factor gene reduces the activity of the molecule by 50%. That is, the carboxyl-terminal region is

particularly sensitive to substitution with a homologous sequence from a heterologous protein, such a substitution having sufficient impact on protein function to decrease activity by 50%. A similar decrease in activity is seen following substitution of the amino terminus (Suter, et al., 1992).

All Ly-6 terminal CIPs are believed to share the property of being attached to cell membranes by means of a GPI linkage. As understood in the art, the addition of such a GPI moiety to a nascent protein coincides with a proteolytic processing step that removes a number of amino acid residues from the carboxyl-terminus of the polypeptide. Accordingly, mature Ly-6 terminal CIPs do not include all of the amino acids specified by the full length nucleic acid molecules that encode them. Specifically, they do not include some or all of the amino acid residues downstream of the cysteine backbone Ly-6 motif, e.g., the amino acids downstream of cysteine 69 of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7 (CD59) and downstream of cysteine 72 of SEQ ID NO:3. (As used herein, "downstream" means towards the carboxyl terminus of the polypeptide or towards the 3' end of the coding strand of the nucleic acid molecule coding for the polypeptide and "upstream" means towards the amino terminus of the polypeptide or towards the 5' end of the coding strand of the nucleic acid molecule coding for the polypeptide.) It is not known which amino acids downstream of the Ly-6 cysteine backbone motif are present or absent in any of these terminal CIPs when they are in the mature, GPI anchored state.

As discussed in detail below, the present invention involves the removal of selected amino acids of such Ly-6 terminal CIPs downstream of the Ly-6 motif. In view of the foregoing state of the art, it was not known, prior to the present invention, what effects such amino acid removal would have on terminal CIP function. In

particular, it was not known if Ly-6 terminal CIPs would retain any complement inhibitory activity after such removal.

5 Various attempts have been made to examine the effects of GPI anchors on protein function. In the case of CD55, the substitution of protein fragments that contain a transmembrane domain for the carboxyl-terminal sequences believed to be involved in the addition of the GPI anchor (referred to hereinafter as the "GPI signal
10 sequence") results in a protein with equal activity to the native GPI-anchored protein (Lublin, et al., 1991). In the case of the Ly-6 protein, Ly-6E (Ly-6E.1), which is a GPI-anchored cell surface protein that is structurally related to CD59 (Philbrick, et al. 1990),
15 the substitution of a fragment containing a transmembrane domain for the carboxyl-terminal GPI signal sequences downstream of the Ly-6 motif produces a non-functional protein, i.e., a protein which is not capable of activating T-cells (Su, et al., 1991).

20 SUMMARY OF THE INVENTION

In view of the foregoing, it is an object of this invention to provide novel proteins that can be used in controlling the complement system of humans and other animals. It is a further object of the invention to
25 provide nucleic acid sequences and associated genetic engineering constructs for producing such proteins either in vitro or in vivo.

More particularly, it is the object of the invention to provide novel proteins that are Ly-6 terminal
30 complement inhibitors, but are anchored to the cell surface by means independent of GPI anchoring. It is an additional object of the invention to provide molecules of this type that will not transmit an activating signal into the cells to which they are bound, e.g., endothelial
35 cells, lymphocytes, or platelets, either after antibody crosslinking, or upon binding of the terminal CIP to its ligand. It is a further object of the invention to

provide molecules of this type that cannot be removed from the surfaces of the cells to which they are bound by the actions of lipid cleaving enzymes such as phospholipases and which are not preferentially incorporated into shed vesicles.

To achieve the foregoing and other objects, the present invention, in accordance with certain of its aspects, provides the complete cDNA sequences of chimeric genes encoding chimeric protein products which comprise the fusion of a Ly-6 terminal CIP with a heterologous transmembrane (TM) domain. Prior to fusion, selected amino acid residues located downstream from the Ly-6 motif of the terminal CIP are deleted. The invention also comprises the chimeric protein products encoded by these genes, such chimeric molecules being referred to hereinafter as TMTCIPs (i.e., transmembrane terminal complement inhibitor proteins). In the preferred embodiments of the invention, the chimeric proteins have greater than 50% of the complement inhibitory activity of the native, GPI-anchored terminal CIP from which the TMTCIP is derived where such activity is preferably measured using a dye release assay of the type described below in Example 4.

The protection from complement attack offered by the TMTCIPs of the invention can be provided via gene transfer for the therapeutic prevention of pathologic complement attack in, for example, transplantation. In a preferred form of such therapy, the expression of the TMTCIP can be directed to the surfaces of cells of non-human animal organs, e.g., organs of non-human transgenic animals, in order to protect such organs from complement attack upon transplantation into a human patient.

The accompanying drawings, which are incorporated in and constitute part of the specification, illustrate certain aspects of the preferred embodiments of the invention and, together with the description, serve to

explain certain principles of the invention. It is to be understood, of course, that both the drawings and the description are explanatory only and are not restrictive of the invention.

5 BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 shows aligned amino acid sequences of Human, African Green Monkey, Baboon, Owl Monkey, Marmoset, Squirrel Monkey, and Herpesvirus Saimiri Ly-6 terminal CIPs (CD59, AGMCIP, BABCIP, OWMCIP, MARCIP, SQMCIP, and HVS-15, respectively). The cysteine residues making up the Ly-6 cysteine backbone motif of each protein are underlined.

FIG. 2 shows a comparison of the cell surface expression of CD59 epitopes on Balb/3T3 cells. The three traces represent cell surface expression profiles of a positive Balb/3T3 clone expressing the CD59-MCP TMTCIP (CD59-TM), a native human CD59 transfectant (CD59-GPI) as a positive control and a vector (pcDNA3, Invitrogen, San Diego, CA) without insert transfectant (Vector Control) as a negative control.

FIG. 3 shows a comparison of the cell surface expression of CD59 epitopes on mouse L cells. The broad trace represents cell surface expression profiles of pooled L cells transduced with retroviral virion particles generated using the pL-CD59-MCP-TM-SN vector (CD59-TM). Also shown are profiles of pooled L cells transduced either with retroviral virion particles generated using the pL-CD59-GPI-SN vector (CD59-GPI) or with retroviral virion particles generated using the pLXSN vector with no insert (Vector Control), as negative controls.

FIG. 4 shows cell surface levels of CD59 antigens on stably transfected Balb/3T3 cells before and after PI-PLC digestion. FIG. 4A shows data obtained using a clone expressing the native human CD59 molecule (CD59-GPI). FIG. 4B shows data obtained using a clone expressing the CD59-MCP TMTCIP (CD59-TM). In each panel, the traces

labeled "A" and "B" represent cells stained with the secondary antibody alone, without or with PI-PLC treatment, respectively. In each panel, the traces labeled "C" and "D" represent cells stained with both the primary (CD59 specific) antibody and the secondary antibody with or without PI-PLC treatment, respectively.

FIG. 5 shows data obtained from dye release assays performed using the transfected Balb/3T3 cells employed in obtaining the data of FIG. 2 and FIG. 4. The cells were challenged with 20% human C8 depleted serum supplemented with a mixture of equal parts of purified human C8 and C9. The amounts, in micrograms per milliliter final concentration, of the mixture of human C8 and C9 added are indicated on the abscissa and the percent of dye release is indicated on the ordinate.

FIG. 6 shows data obtained from dye release assays performed using the transfected mouse L cells employed in obtaining the data of FIG. 3. The cells were challenged with 20% human C8 depleted serum supplemented with a mixture of equal parts of purified human C8 and C9. The amounts, in micrograms per milliliter final concentration, of the mixture of human C8 and C9 added are indicated on the abscissa and the percent of dye release is indicated on the ordinate.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

As discussed above, the present invention provides the complete cDNA sequences of chimeric genes encoding chimeric protein products which comprise the fusion of a Ly-6 terminal CIP and a heterologous transmembrane domain.

I. Terminal CIPs

A variety of terminal CIPs can be used in the practice of the invention. In particular, Ly-6 terminal CIPs can be used. In addition to sharing the homologies shown in formulas (1) and (2) above, the Ly-6 terminal CIPs also share a variety of other homologies which can be seen in the aligned amino acid sequences of FIG. 1.

Homologies found downstream from the Ly-6 cysteine motif of these terminal CIPs include an N immediately following the last C of the Ly-6 motif, another N 6-8 residues downstream from the last C of the Ly-6 motif (referred to herein as the "truncation-Asn"), and the following consensus sequence, hereinafter referred to as the "downstream consensus sequence" which includes the aforementioned truncation-Asn at the third position in the sequence:

10 (L or I) (E or K) N (G or I) (G or K) (T or R) (S or T) (L or I) S (K or E or D) K (T or A) (V or I or L) (L or V) L L (V or L) (A or T or I) (P or L) (F or L) L (A or V) (A or T) A W (S or C or N) (L or R or F) (H or P) (P or L).

15 In addition to these structural commonalities, testing of the Ly-6 terminal CIPs of FIG. 1 has shown that they share the ability to substantially inhibit the activity of human complement. (See U.S. patent application Serial No. 08/105,735, and PCT patent application Serial No. PCT/US93/00672, referred to above.) In particular, each of CD59, AGMCIP, BABCIP, OWMCIP, SQMCIP, and HVS-15 had substantial human complement inhibitory activity. MARCIP was not tested, but is also expected to have such activity.

25 II. Transmembrane Domains

As known in the art, transmembrane proteins may span the membrane once or several times along the length of their amino acid chains. There are in general two different ways in which a transmembrane protein that spans the membrane only once may be embedded in a membrane. Most commonly, these proteins have their single transmembrane domain located towards the carboxyl-terminal end of the polypeptide chain and are oriented so that the region amino-terminal to the transmembrane domain is outside the cell or in a non-cytoplasmic cellular compartment and the region carboxyl-terminal to the transmembrane domain is in the

30

35

cytoplasmic compartment. The second orientation of a transmembrane protein with a single membrane spanning transmembrane domain is the opposite of this common arrangement, that is, the region amino-terminal to the transmembrane domain is in the cytoplasmic compartment and the region carboxyl-terminal to the transmembrane domain is located outside the cell or in a non-cytoplasmic cellular compartment.

Other transmembrane proteins cross the membrane several times. Most commonly, eukaryotic representatives of this type of transmembrane protein have seven consecutive transmembrane domains, most of them connected by short hydrophilic loop regions.

Transmembrane proteins in general include at least one contiguous stretch of amino acid residues which resides in the lipid bilayer membrane (referred to hereinafter as "membrane amino acids"), and at least two contiguous stretches of amino acid residues which extend away from the membrane, one generally cytoplasmic (referred to hereinafter as "cytoplasmic amino acids"), and one generally extracellular or sequestered in a non-cytoplasmic cellular compartment (referred to hereinafter as "extracellular amino acids"). As referred to herein, cytoplasmic amino acids and extracellular amino acids always include at least one charged amino acid residue immediately adjacent to the membrane amino acids (referred to herein as the "first cytoplasmic amino acid" and the "first extracellular amino acid," respectively).

Membrane amino acids are characterized as groups of at least about 20 amino acids (the minimum generally needed to span a membrane), most of which are hydrophobic (uncharged) amino acids. Charged (hydrophilic) amino acids are usually absent from these groups, but in some cases two hydrophilic residues of opposite charge may lie close together inside the membrane where they neutralize each other.

-22-

Transmembrane domains derived from a variety of transmembrane proteins can be used in the practice of the invention. However, transmembrane domains with cytoplasmic amino acids which include cysteine residues in close proximity to the first cytoplasmic amino acid may be expressed at lower levels on the cell surface than transmembrane domains that do not contain such cysteines. This decreased expression is believed to result from the propensity of these cysteine residues to form intermolecular bonds with similarly placed cysteines of adjacent nascent transmembrane protein molecules. Such intermolecular cysteine linkages cause aggregation of the nascent transmembrane proteins, generally within the Golgi apparatus (where newly synthesized transmembrane proteins are processed within the typical cell), and thus block the transport of such nascent proteins to the cell surface.

With regard to the TMTCIP molecules of the invention, it is notable that transfection of mammalian cells with an expression vector encoding a chimeric terminal CIP containing a putative transmembrane domain from the herpesvirus saimiri CAPH gene (see PCT patent application Serial No. PCT/US93/00672, mentioned above) does not result in high enough levels of cell surface expression of terminal CIP epitopes to be detected by FACS analysis (see Example 1). The cytoplasmic amino acids of this putative transmembrane domain include cysteine residues spaced two and five amino acids from the first cytoplasmic amino acid, a histidine. The presence of these cysteines is believed to be responsible for the low levels of expression seen with this putative transmembrane domain. For this reason, transmembrane domains of this type are not preferred for use with the present invention.

As used herein, the term "transmembrane domain" is intended to comprise: 1) the portion of a transmembrane protein which spans the membrane, i.e., the at least

about twenty membrane amino acids normally required for this purpose, 2) the adjacent predominantly charged cytoplasmic amino acids within about five to about ten residues from the membrane amino acids, and 3) the adjacent predominantly charged extracellular amino acids within about five to about ten residues from the membrane amino acids. These adjacent predominantly charged cytoplasmic and extracellular amino acids are involved in anchoring the protein in the membrane. As discussed above, preferred transmembrane domains do not include cytoplasmic amino acids that are cysteine residues within five amino acids of the first cytoplasmic amino acid.

While it is possible to examine a protein sequence and pick out a region with about 20 consecutive hydrophobic amino acids, some transmembrane domains, as discussed above, contain a small number of hydrophilic amino acids interspersed within their predominantly hydrophobic residues. Accordingly, transmembrane domains are more effectively identified by using hydrophobicity scales to compute hydropathy plots (Branden, et al., 1991).

Hydrophobicity scales provide a numerical value for the hydrophobicity of individual amino acids. These scales have been developed on the basis of solubility measurements of amino acids in different solvents, vapor pressures of side-chain analogues, analysis of side-chain distributions within soluble proteins, and theoretical energy calculations (Kyte, et al., 1982; and Engelman, et al., 1986).

Hydropathy plots are computed from amino acid sequences using hydrophobicity values as follows. First, for each position in the sequence, a hydropathic index is calculated. The hydropathic index is the mean value of the hydrophobicity of the amino acids within a "window," usually 19 residues long, around each position. The hydropathic indices are then plotted versus amino acid sequence position to produce the hydropathy plot.

Transmembrane domains are then identified from the hydropathy plots by searching for regions where the hydropathic index is high for a number of consecutive positions in the sequence, e.g., by searching for regions with broad peaks with high positive (i.e., hydrophobic) values.

In terms of the present invention, the transmembrane domain will preferably have a hydropathic index greater than about +0.5, using the scale of Kyte et al. (Kyte et al., 1982) and a window of 19 amino acids, over a region of at least about 12 amino acid residues.

Additional contiguous amino acids of the transmembrane protein can be included in or encoded by the chimeric molecules of the invention provided those additional amino acids do not substantially impair the insertion of the transmembrane domain into the membrane, the transport of the nascent chimeric protein to the cell surface, or the complement inhibitory activity of the terminal CIP portion of the chimeric molecule.

While the molecules of the present invention may be constructed with any functional transmembrane domain, one derived from a protein with only a single transmembrane domain and having the region carboxyl-terminal to its transmembrane domain in the cytoplasm is preferred. A large number of such proteins have been reported in the literature, including the following: CD46; the major histocompatibility antigens and related transmembrane proteins of the immunoglobulin multigene superfamily including intercellular adhesion molecules, such as ICAM-1 (CD54), ICAM-2, ICAM-3, VCAM-1, PECAM-1 (CD31) and HCAM (CD44); the selectins, including E-selectin, L-selectin, and P-selectin (CD62); the Alzheimer's amyloid precursor protein; the insulin receptor; the epidermal growth factor receptor; the gp41 protein of the AIDS virus, HIV; the p21 proteins of HTLV1 and HTLV2; and the p15E proteins of the murine and feline leukemia viruses.

TM domains derived from any of these proteins, as well as from other transmembrane proteins, can be used in the practice of the invention. These domains can be most easily used by incorporating into the chimeric molecule the entire carboxyl end of the transmembrane protein beginning at a point upstream from the transmembrane domain. A particularly preferred TM domain is that constituting amino acids 294 to 326 of CD46 (MCP, SEQ ID NO:8). This domain can be conveniently used along with amino acids 327 to 350, which comprise the carboxyl end of the CD46 protein downstream from the transmembrane domain of this molecule, and along with amino acids 270 to 293 upstream of the TM domain which do not interfere with insertion of the CD46 TM domain into cell membranes and, as shown below, do not inhibit the complement inhibitory activity of Ly-6 terminal CIPs.

In addition to using hydropathy plots to identify TM domains suitable for use in the present invention, such domains can also be identified biochemically using, for example, protease digestion techniques or by making chimeric molecules containing soluble proteins operatively linked to signal sequences and containing putative transmembrane domains, and assaying for membrane insertion of the chimeric protein.

25 III. TMTCIP Genes and Vectors Containing Such Genes

The isolation, truncation, and fusion of the nucleic acid fragments encoding the terminal CIP and the TM domain are performed using recombinant nucleic acid techniques known in the art, including: PCR generation of the desired fragments and/or restriction digestion of cloned genes; PCR fusion of the desired fragments; or enzymatic ligation of restriction digestion products (Sambrook, et al., 1989; and Ausubel et al., 1992). Alternatively, the nucleic acid molecules encoding the TMTCIPs of the invention or any or all of the nucleic acid fragments used to assemble the chimeric genes for

the TMTCIPs can be synthesized by chemical means (Talib, et al., 1991).

The chimeric genes of the invention are prepared by 1) truncating the nucleic acid sequence for a Ly-6 terminal CIP so as to remove selected amino acid residues downstream of the Ly-6 motif in order to inactivate the normal GPI signal sequence, and 2) fusing the truncated sequence to a sequence coding for a selected TM domain and desired amino acids surrounding the TM domain.

The truncation of the nucleic acid sequence encoding the Ly-6 terminal CIP will remove at least some of the carboxyl-terminal amino acid residues downstream from the Asn which is located between 6 and 8 amino acid residues after the last (tenth) Cys of the Ly-6 motif. This Asn is also located at the third position in the downstream consensus sequence presented above, i.e., it is the truncation-Asn defined above. All known Ly-6 terminal CIPs include such a truncation-Asn.

In some cases, all of the amino acid residues after the truncation-Asn are removed. Alternatively, less than all can be removed, the criterion being that sufficient numbers of residues are removed so that the GPI signal sequence is inoperative. In general, the simplest approach is to remove all amino acid residues downstream of the truncation-Asn. If desired, the truncation can extend further upstream from the truncation-Asn, preferably starting at a point downstream from the last Cys of the Ly-6 motif. Truncations beginning upstream from the last Cys of the Ly-6 motif are in general not preferred, but can be used if desired. The criterion for truncations upstream of the truncation-Asn is the requirement that the TMTCIP has greater than 50% of the complement inhibitory activity of the parent (native) Ly-6 terminal CIP.

In terms of the Ly-6 terminal CIPs of FIG. 1, the preferred truncation comprises all of the amino acids downstream of Asn 77 of BABCIP (SEQ ID NO:1), Asn 75 of

AGMCIP (SEQ ID NO:2), Asn 80 of SQMCIP (SEQ ID NO:3), Asn 77 of OWMCIP (SEQ ID NO:4), Asn 77 of MARCIP (SEQ ID NO:5), Asn 77 of HVS-15 (SEQ ID NO:6), and Asn 77 of CD59 (SEQ ID NO:7). Of these Ly-6 terminal CIPs, CD59 is preferred. As discussed above, a preferred TM domain is from CD46. Accordingly, a particularly preferred embodiment of the invention comprises residues 1-77 of CD59 (SEQ ID NO:7) fused to amino acids 270-350 of CD46 (SEQ ID NO:8)

In addition to the foregoing, the present invention provides recombinant expression vectors which include nucleic acid fragments encoding the chimeric TMTCIPs of the invention. The nucleic acid molecule coding for such a chimeric protein can be inserted into an appropriate expression vector, i.e., a vector that contains the necessary elements for the transcription and translation of the inserted protein-encoding sequence. The necessary transcriptional and translational signals can also be supplied by the genes used to construct the fusion genes of the invention and/or their flanking regions.

The transcriptional and translational control sequences for expression vector systems to be used to direct expression in vertebrate cells may be provided by viral sources. For example, commonly used promoters and enhancers are derived from Polyoma virus, Adenovirus, Simian Virus 40 (SV40), the Molony murine leukemia virus (MMLV), including the long terminal repeat (MMLV-LTR), and human cytomegalovirus (CMV), including the cytomegalovirus immediate-early gene 1 promoter and enhancer. Retroviral expression vectors are a preferred system for expression of the TMTCIPs of the invention.

The manipulation of retroviral nucleic acids to construct retroviral vectors and packaging cells is accomplished using techniques known in the art. See Ausubel, et al., 1992, Volume 1, Section III (units 9.10.1 - 9.14.3); Sambrook, et al., 1989; Miller, et al., 1989; Eglitis, et al., 1988; U.S. Patents Nos. 4,650,764,

4,861,719, 4,980,289, 5,122,767, and 5,124,263; as well as PCT Patent Publications Nos. WO 85/05629, WO 89/07150, WO 90/02797, WO 90/02806, WO 90/13641, WO 92/05266, WO 92/07943, WO 92/14829, and WO 93/14188.

5 In particular, the retroviral vectors of the invention can be prepared and used as follows. First, a TMTCIP retroviral vector is constructed and packaged into non-infectious transducing viral particles (virions) using an amphotropic packaging system, preferably one
10 suitable for use in gene therapy applications.

Examples of such packaging systems are found in, for example, Miller, et al., 1986; Markowitz, et al., 1988; Cosset, et al., 1990; U.S. Patents Nos. 4,650,764, 4,861,719, 4,980,289, 5,122,767, and 5,124,263, and PCT
15 Patent Publications Nos. WO 85/05629, WO 89/07150, WO 90/02797, WO 90/02806, WO 90/13641, WO 92/05266, WO 92/07943, WO 92/14829, and WO 93/14188. A preferred packaging cell is the PA317 packaging cell line (ATCC CRL 9078).

20 The generation of "producer cells" is accomplished by introducing retroviral vectors into the packaging cells. Examples of such retroviral vectors are found in, for example, Korman, et al., 1987; Morgenstern, et al., 1990; U.S. Patents Nos. 4,405,712, 4,980,289, and
25 5,112,767; and PCT Patent Publications Nos. WO 85/05629, WO 90/02797, and WO 92/07943. A preferred retroviral vector is the MMLV derived expression vector pLXSN (Miller, et al., 1989). The retroviral vector used in the practice of the present invention will be modified to
30 include the chimeric gene encoding the TMTCIP.

The producer cells generated by the foregoing procedures are used to produce the retroviral vector particles (virions). This is accomplished by culturing of the cells in a suitable growth medium. Preferably,
35 the virions are harvested from the culture and administered to the target cells which are to be transduced, e.g., xenogeneic cells to be used for

transplantation into a patient whose complement can be inhibited by the Ly-6 terminal CIP of the TMT-CIP, cells of a xenogeneic organ to be used for transplantation into such a patient, the patient's own cells, and other cells to be protected from complement attack, as well as stem cells such as embryonic stem cells, which can be used to generate transgenic cells, tissues, or organs for transplantation. Alternatively, when practicable, the target cells can be co-cultured with the producer cells. Suitable buffers and conditions for stable storage and subsequent use of the virions can be found in, for example, Ausubel, et al., 1992.

Pharmaceutical compositions containing the retroviral vector particles of the invention can be administered in a variety of unit dosage forms. The dose will vary according to, e.g., the particular vector, the manner of administration, the particular disease being treated and its severity, the overall health and condition and age of the patient, the condition of the cells being treated, and the judgment of the physician. Dosage levels for transduction of mammalian cells are generally between about 10^6 and 10^{14} colony forming units of retroviral vector particles per treatment.

A variety of pharmaceutical formulations can be used for administration of the retroviral vector particles of the invention. Suitable formulations are found in, for example, Remington's Pharmaceutical Sciences, Mack Publishing Company, Philadelphia, PA, 17th ed., 1985, and will include a pharmaceutically effective carrier, such as saline, buffered (e.g., phosphate buffered) saline, Hank's solution, Ringer's solution, dextrose/saline, glucose solutions, and the like. The formulations may contain pharmaceutically acceptable auxiliary substances as required, such as, tonicity adjusting agents, wetting agents, bactericidal agents, preservatives, stabilizers, and the like.

IV. Transgenic Animals

In accordance with certain aspects of the invention, the nucleic acid molecules of the present invention are used to generate engineered transgenic animals (for example, rodent, e.g., mouse, rat, capybara, and the like, lagomorph, e.g., rabbit, hare, and the like, ungulate, e.g., pig, cow, goat, sheep, and the like, etc.) that express the TMTCIPs of the invention on the surfaces of their cells (e.g., endothelial cells) using techniques known in the art. These techniques include, but are not limited to, microinjection, e.g., of pronuclei, electroporation of ova or zygotes, nuclear transplantation, and/or the stable transfection or transduction of embryonic stem cells derived from the animal of choice.

A common element of these techniques involves the preparation of a transgene transcription unit. Such a unit comprises a DNA molecule which generally includes: 1) a promoter, 2) the nucleic acid sequence of interest, i.e., the sequence encoding the TMTCIP of the present invention, and 3) a polyadenylation signal sequence. Other sequences, such as, enhancer and intron sequences, can be included if desired. The unit can be conveniently prepared by isolating a restriction fragment of a plasmid vector which expresses the TMTCIP protein in, for example, mammalian cells. Preferably, the restriction fragment is free of sequences which direct replication in bacterial host cells since such sequences are known to have deleterious effects on embryo viability.

The most well known method for making transgenic animals is that used to produce transgenic mice by superovulation of a donor female, surgical removal of the egg, injection of the transgene transcription unit into the pro-nuclei of the embryo, and introduction of the transgenic embryo into the reproductive tract of a pseudopregnant host mother, usually of the same species. See Wagner, U.S. Patent No. 4,873,191, Brinster, et al.,

1985, Hogan, et al., 1986, Robertson 1987, Pedersen, et al., 1990.

The use of this method to make transgenic livestock is also widely practiced by those of skill in the art. As an example, transgenic swine are routinely produced by the microinjection of a transgene transcription unit into pig embryos. See, for example, PCT Publication No. WO92/11757 In brief, this procedure may, for example, be performed as follows.

First, the transgene transcription unit is gel isolated and extensively purified through, for example, an ELUTIP column (Schleicher & Schuell, Keene, NH), dialyzed against pyrogen free injection buffer (10mM Tris, pH7.4 + 0.1mM EDTA in pyrogen free water) and used for embryo injection.

Embryos are recovered from the oviduct of a hormonally synchronized, ovulation induced sow, preferably at the pronuclear stage. They are placed into a 1.5 ml microfuge tube containing approximately 0.5 ml of embryo transfer media (phosphate buffered saline with 10% fetal calf serum). These are centrifuged for 12 minutes at 16,000 x g in a microcentrifuge. Embryos are removed from the microfuge tube with a drawn and polished Pasteur pipette and placed into a 35 mm petri dish for examination. If the cytoplasm is still opaque with lipid such that the pronuclei are not clearly visible, the embryos are centrifuged again for an additional 15 minutes. Embryos to be microinjected are placed into a drop of media (approximately 100 μ l) in the center of the lid of a 100 mm petri dish. Silicone oil is used to cover this drop and to fill the lid to prevent the medium from evaporating. The petri dish lid containing the embryos is set onto an inverted microscope equipped with both a heated stage (37.5-38°C) and Hoffman modulation contrast optics (200X final magnification). A finely drawn and polished micropipette is used to stabilize the embryos while about 1-2 picoliters of injection buffer

containing approximately 200-500 copies of the purified transgene transcription unit is delivered into the nucleus, preferably the male pronucleus, with another finely drawn and polished micropipette. Embryos
5 surviving the microinjection process as judged by morphological observation are loaded into a polypropylene tube (2 mm ID) for transfer into the recipient pseudopregnant sow.

Offspring are tested for the presence of the
10 transgene by isolating genomic DNA from tissue removed from the tail of each piglet and subjecting about 5 micrograms of this genomic DNA to nucleic acid hybridization analysis with a transgene specific probe.

Another commonly used technique for generating
15 transgenic animals involves the genetic manipulation of embryonic stem cells (ES cells) as described in PCT Patent Publication No. WO 93/02188 and Robertson, 1987.

In accordance with this technique, ES cells are grown as described in, for example, Robertson, 1987, and in U.S.
20 Patent No. 5,166,065 to Williams et al. Genetic material is introduced into the embryonic stem cells by, for example, electroporation according, for example, to the method of McMahon, et al., 1990, or by transduction with a retroviral vector according, for example, to the method
25 of Robertson, et al., 1986, or by any of the various techniques described by Lovell-Badge, 1987.

Chimeric animals are generated as described, for example, in Bradley, 1987. Briefly, genetically modified ES cells are introduced into blastocysts and the modified
30 blastocysts are then implanted in pseudo-pregnant female animals. Chimeras are selected from the offspring, for example by the observation of mosaic coat coloration resulting from differences in the strain used to prepare the ES cells and the strain used to prepare the
35 blastocysts, and are bred to produce non-chimeric transgenic animals.

- 33 -

Other methods for the production of transgenic animals are disclosed in U.S. Patent No. 5,032,407 to Wagner et al., and PCT Publication No. WO90/08832.

Among other applications, transgenic animals prepared in accordance with the invention are useful as model systems for testing the xenotransplantation of their engineered tissues or organs and as sources of engineered tissues or organs for xenotransplantation. The expression of functional TMTCIPs on the surfaces of endothelial cells and/or other cell types in the tissues and organs (e.g., hormone producing cells such as those in the pancreatic islets) of the transgenic animals will provide enhanced protection to those cells, tissues and organs from hyperacute complement-mediated rejection following xenotransplantation in recipient animals, e.g., humans, whose complement can be inhibited by the Ly-6 terminal CIP of the TMTCIP. In addition to their use in producing organs for transplantation, the TMTCIP nucleic acid constructs of the invention can also be used to engineer cultured cells (e.g., endothelial cells) of various species for subsequent use in transplantation.

V. Representative Modifications

Although specific embodiments of the invention are described and illustrated herein, it is to be understood that modifications can be made without departing from the invention's spirit and scope.

For example, the primary amino acid structures of the TMTCIPs of the invention may be modified by creating amino acid substitutions or nucleic acid mutations. At least some complement regulatory activity should remain after such modifications. Similarly, nucleic acid mutations which do not change the amino acid sequences, e.g., third nucleotide changes in degenerate codons, are included within the scope of the invention. Also included are sequences comprising changes that are found as naturally occurring allelic variants of the CIP and TM genes used to create the TMTCIPs.

Without intending to limit it in any manner, the present invention will be more fully described by the following examples.

Example 1

5 Expression Vectors and Retroviral Virion Particles
 Comprising a CD59/MCP TMTCIP

 A transmembrane form of CD59 (CD59-TM) was constructed according to the present invention by replacing the carboxyl-terminal region containing the
10 GPI- anchor signal of CD59 with the carboxyl-terminal region, including the transmembrane domain, of MCP (CD46). An approximately 314 bp restriction fragment (hereinafter referred to as CD59₇₇) containing CD59 truncated at the "truncation-Asn" described above, i.e.,
15 amino acid 77 of the mature protein, was prepared by digestion of plasmid pCD59/CCPH (see below) with SspI and BamHI.

 The carboxyl-terminus of CD46 was PCR amplified using HeLa cell reverse-transcribed mRNA as template and
20 the following primers: 5'-CGCGAGGCCT ACTTACAAGC CTCCAG-3' (SEQ ID NO:9) and 5'-CGCGCTATTC AGCCTCTCTG CTCTGC-3' (SEQ ID NO:10). These oligonucleotides amplified a fragment coding for amino acids 270-350 of the mature CD46 protein, a region shown previously to
25 comprise a functional transmembrane domain (Lublin, et al., 1991). The approximately 250 bp fragment produced by this PCR reaction was cloned into a plasmid vector using the T/A cloning kit (Invitrogen, San Diego, CA). The pCRII plasmid vector included in this kit served as
30 the recipient, and the resulting plasmid construct was amplified in E. coli and purified. The MCP insert was subsequently sequenced to confirm that the plasmid contained the sequence shown in SEQ ID NO:11.

 An endogenous StuI site found at the 5' end of the
35 CD46 PCR fragment was utilized to ligate this domain to the SspI site at the 3' end of CD59₇₇ in the eukaryotic expression vector pcDNA3 (Invitrogen, San Diego, CA) to

-35-

yield plasmid pcDNA3/CD59-MCP-TM (ATCC designation 69530).

5 The resulting construct was linearized with EcoRI, the unpaired ends were filled, and BamHI linkers (#1071, New England Biolabs, Tozer, MA) were ligated onto the resulting blunt ends. This linkered construct was digested with BamHI and the liberated fragment was subcloned into the BamHI site of the retroviral vector pLXSN (Miller, et al., 1989) to yield pL-CD59-MCP-TM-SN.

10 Constructs with the correct orientation for expression were identified by restriction enzyme analysis and confirmed by sequencing.

A DNA fragment encoding the carboxyl-terminus of the CCPH gene was prepared by PCR amplification using plasmid pKS-/mCCPH (ATCC designation 69178) as template and the following primers: 5'-CCGGACCTGT GTAACCTTTAA CGAACAGCTT GAAAATATTG GTAGGATATG CAATGGAAAT TGTTACAAC-3' (SEQ ID NO:12) and 5'-TAGTTACTGC CCGGACATGC-3' (SEQ ID NO:13).

20 As described above for the MCP PCR fragment, the approximately 250 bp CCPH PCR product was cloned into plasmid pCRII, yielding plasmid pCRII/CCPH, and the CCPH insert was sequenced to confirm that the plasmid contained the desired sequence, in this case SEQ ID NO:14.

25 The pCRII/CCPH plasmid was then digested with AvaII and EcoRI, and the insert fragment was purified and subcloned in a three-way ligation reaction with plasmid pcDNA/AMP (Invitrogen) cut with BamHI and EcoRI and an approximately 300 base pair BamHI - AvaII fragment

30 isolated from a full length CD59 cDNA construct in pUC19 (Philbrick et al., 1990). The product of this three-way ligation is referred to herein as plasmid pCD59/CCPH.

This plasmid was transfected into Balb/3T3 cells and the cells were assayed for cell surface expression of

35 CD59 epitopes by indirect immunofluorescence as described below in Example 2. As discussed above, the putative TM domain of CCPH contains two cytoplasmic amino acids,

within five amino acids of the first cytoplasmic amino acid, that are cysteine residues, a characteristic that is believed to result in low levels of cell surface expression. Cell surface expression of CD59 epitopes was indeed below the levels detectable by the indirect immunofluorescence assay.

Control vectors.

Full-length CD59 containing the GPI-anchor signal (CD59-GPI) was cloned into BamHI - EcoRI digested pcDNA3 (Invitrogen) as an BamHI - EcoRI fragment obtained from plasmid pc8-hCD59-103 (ATCC designation 69231) to yield plasmid pcDNA3-CD59-GPI.

Retroviral vector plasmid pL-CD59-GPI-SN was produced by isolating an approximately 1100 bp EcoRI fragment from a full length CD59 cDNA construct in pUC19 (Philbrick et al., 1990) and ligating this fragment into plasmid pLXSN. Constructs with the correct orientation for expression were identified by restriction enzyme analysis.

Amphotropic virus production.

Amphotropic virus was produced through an intermediate ecotropic packaging cell line as described in Warren et al., 1987. Briefly, psi 2 cells (obtained from Dr. Stephen L. Warren, Department of Pathology, Yale University School of Medicine, New Haven, CT) were transfected with pLXSN or the pLXSN constructs described above, i.e., pL-CD59-MCP-TM-SN or pL-CD59-GPI-SN, using DMSO shock followed by selection in DMEM containing 500 µg/ml (active) G418 and 10% heat inactivated FCS. Transfectants were pooled and a 24 hour supernatant was harvested from the cells at 90% confluency. The ecotropic virus stock was used to infect the amphotropic packaging cell line PA317 (ATCC designation CRL 9078). These cells were also selected in the same medium with G418 following which a virus stock was collected from pooled transductants in the same medium without G418.

Example 2Expression of the CD59/MCP TMTCIP by Mammalian Cells

Cells of the murine fibroblast cell line, Balb/3T3 (ATCC designation CCL 163) were stably transfected with
5 pcDNA3-CD59-GPI, pcDNA3/CD59-MCP-TM, or pcDNA3 alone using the calcium phosphate method (Ausubel, et al., 1992). Cells were selected in DMEM containing 10% heat inactivated FCS and 500 $\mu\text{g/ml}$ of G418 (active) and colonies were isolated using cloning cylinders.

10 Mouse L cells were obtained from Dr. Peter Cresswell, Immunobiology Department, Yale University School of Medicine, New Haven, CT. Such mouse L cells are unable to express GPI anchored proteins (Ferguson, et al., 1988). L cells were transduced with the amphotropic virus supernatants obtained using pL-CD59-GPI-SN,
15 pL-CD59-MCP-TM-SN, or pLXSN alone by adding 1 ml of the virus stock to 5×10^5 L cells in medium containing 8 $\mu\text{g/ml}$ polybrene. After an overnight incubation, medium containing 500 $\mu\text{g/ml}$ G418 was added and selection
20 continued for 14 days. Transduced L cells were selected and analyzed as a pool. G418 resistant cells were assayed for the presence of CD59 antigens on the cell surface by indirect immunofluorescence using monoclonal and polyclonal antibody preparations. A rabbit polyclonal
25 anti-CD59 antibody preparation, #349, that was produced by injecting rabbits with CD59 purified from human erythrocytes as described by Sims et al., 1989, was provided by Dr. Peter Sims (Blood Research Institute, Milwaukee, WI). The anti-CD59 mAb, MEM-43, was
30 purchased from Biodesign International, Kennebunkport, ME.

Cell surface indirect immunofluorescence analysis was typically performed on 2.5×10^5 cells with 50 $\mu\text{g/ml}$ of the primary polyclonal antibody or 20 $\mu\text{g/ml}$ of the
35 monoclonal antibody in 1xPBS containing 2% fetal bovine serum. Goat anti-rabbit IgG or goat anti-mouse IgG FITC conjugated antisera were used as secondary antibodies

-38-

(Zymed Laboratories, South San Francisco, CA). Fluorescence was measured using a FACSsort instrument (Becton-Dickinson Immunocytometry Systems, San Jose, CA).

5 FIG. 2 illustrates cell surface expression profiles of a positive Balb/3T3 clone expressing CD59-TM, as well as of a native human CD59 (CD59-GPI) transfectant as a positive control and a vector (pcDNA3) without insert (vector control) transfectant as a negative control. As
10 shown therein, essentially the same amount of anti-CD59 antibody bound to the surfaces of cells expressing the CD59-TM fusion protein as bound to the positive control cells expressing native CD59. This result shows that
15 equivalent amounts of CD59 antigens were present on the Balb/3T3 cells of the invention (CD59-TM) and those of the positive control (CD59-GPI).

 The pooled L cell transfectants showed a wide range of CD59-TM expression while, as expected in cells that cannot express GPI anchored proteins, CD59-GPI was not
20 expressed (Figure 3).

Example 3

TMTCIP Expressed in Mammalian Cells Is Not Affected By Phosphatidylinositol-Phospholipase C Digestion

 To test for the presence of a GPI anchor, cells were
25 treated with phosphatidylinositol-phospholipase C (PI-PLC, Boehringer-Mannheim Corporation, Biomedical Products Division, Indianapolis, Indiana) at 1 U/ml for 1 hr at 37°C prior to FACS analysis. This treatment hydrolyzes (cleaves) GPI anchors, and thus frees GPI
30 anchored proteins from the cell surface. PI-PLC digestion was performed on Balb/3T3 cells expressing the CD59-TM TMTCIP (or CD59-GPI as a control). The results of these experiments are presented in Figure 4. In these experiments, mock treated cells (no PI-PLC) retained the
35 TMTCIP and native CD59 on their cell surfaces (see curve D in FIG. 4A and FIG. 4B), whereas PI-PLC treatment resulted in the loss of cell surface CIPs from the native

CD59 control cells (see curve C in FIG. 4A), but not the CD59-TM cells (see curve C in FIG. 4B). These experiments demonstrate that CD59-TM is not anchored to the cell membrane through a GPI linkage and that CD59-TM is substantially resistant to the action of lipase enzymes which can cleave a glycosyl- phosphatidylinositol (GPI) anchor.

Example 4

Functional Analysis of CD59-TM in Mouse Cells

The functional activity of TMTCIP molecules expressed in transfected mouse Balb/3T3 cells and transduced mouse L cells was assessed by a dye release assay that consisted of measuring the efflux of molecules from the cytoplasm, specifically the cytoplasmic indicator dye, Calcein AM (Molecular Probes, Inc., Eugene, Oregon).

Transfected cells expressing the CD59-TM TMTCIP, as well as cells transfected with the parent expression vectors without CD59-TM encoding inserts (as controls), were grown to confluency in 96-well plates. Cells were washed twice with 200 μ l of Hank's balanced salts solution containing 10 mg/ml bovine serum albumin (HBSS/BSA).

Calcein AM was added (10 μ M final) and the plates were incubated at 37°C for 30 minutes to allow the dye to be internalized by the cells and converted by cellular esterases into a polar fluorescent derivative that is retained inside undamaged cells. The wells were then washed twice with HBSS/BSA to remove dye remaining outside the cells. The cells were then incubated with anti- Balb/3T3 IgG (2 mg/ml in HBSS/BSA), which served as an activator of the classical complement pathway. After a 30 minute incubation at 23°C, unbound IgG was washed away.

The cells were then incubated at 37°C for 30 minutes in the presence of human C8 deficient serum supplemented with purified C8 and C9 to allow complement-mediated

-40-

5 damage to occur. Human C8 depleted serum, as well as purified C8 and C9, were obtained from Quidel Corporation, San Diego, CA. The medium bathing the cells was then transferred to a clean 96-well plate for fluorescence measurement.

10 Under the conditions of this assay, the fluorescent polar derivative of Calcein AM is only released into the medium bathing the test cells if the integrity of the cell membranes is compromised. Therefore, the fluorescence of the Calcein AM released into the medium bathing the test cells versus that retained in the cells provides an indirect, but accurate measure of the level of complement-mediated damage sustained by the cells. Remaining cell-associated dye was determined from a 1% SDS lysate of the cells retained in the 96-well culture plates. This allowed the calculation of percent dye release using the following formulas: Total = released + retained, and, % release = (released / total) x 100. Fluorescence was measured using a Millipore CYTOFLUOR 2350 fluorescence plate reader (490 nm excitation, 530 nm emission).

25 The dye release assays showed that for transfected Balb/3T3 clones expressing equivalent levels of CD59-GPI or CD59-TM (FIG. 2), CD59-TM provided a level of protection from complement attack equivalent to that afforded by the native, GPI-anchored, CD59-GPI molecule (FIG. 5). In particular, cells expressing either of these molecules were approximately 3-fold more effective in preventing complement-mediated lysis at 2.5 µg/ml C8/C9 than cells transfected with the pcDNA3 vector alone, which were readily lysed.

35 These results demonstrate that 1) CD59-TM can be stably expressed on the surface of Balb/3T3 cells, and 2) this chimeric molecule has comparable function to native CD59. The retention of wild-type levels of complement regulatory activity by CD59-TM is of considerable significance in that it shows that the functionality of

the CD59 molecule is not substantially altered by truncation coupled with addition of a TM domain. This result could not have been predicted in advance, especially since other alterations of the CD59 molecule, e.g., truncation of the carboxyl-terminus without addition of a TM domain, or alterations of single amino acids, have been shown to produce molecules with substantially altered expression and/or functionality. See, for example, Nakano, et al., 1993; Norris, et al., 1993; and Petranka, et al., 1993.

Dye release assays were also performed on mouse L cells transduced with the retroviral virion particles generated using the pL-CD59-MCP-TM-SN vector, the pL-CD59- GPI-SN vector, or the no insert pLXSN vector. The results of these experiments are presented in FIG. 6. Only L cells transduced with retroviral particles generated using pL-CD59-MCP-TM-SN demonstrated substantial protection against complement attack. These results demonstrate that the chimeric CD59-TM molecule can successfully be expressed in a cell line unable to express GPI-anchored proteins and that the molecule functions to protect the cells from complement lysis.

The foregoing results show that CD59 retains its Ly-6 terminal complement inhibitor activity when anchored to the cell membrane by a heterologous transmembrane domain, rather than a GPI anchor. This fundamental result, in combination with the conserved nature of all known Ly-6 terminal complement inhibitor proteins (see U.S. patent application Serial No. 08/105,735 and PCT patent application Serial No. PCT/US93/006772), indicates that a heterologous transmembrane domain can be substituted for the GPI signal sequence of a Ly-6 terminal complement inhibitor protein without substantially altering the complement inhibitor activity of the protein.

In comparison to using a native Ly-6 terminal CIP, the TMTCIPs of the invention have the advantages that

-42-

they cannot produce cell activation of the type which depends on the presence of the GPI anchor of the native Ly-6 terminal CIP, and that they cannot be removed from the cell surface by the action of phospholipase enzymes and are less prone to vesicular shedding. These advantages make the TMTCIPs of the invention more suitable than native Ly-6 terminal CIPs for various medical applications, including the facilitation of transplantation of xenogenetic organs.

Although preferred and other embodiments of the invention have been described herein, other embodiments, including a variety of modifications may be perceived and practiced by those skilled in the art without departing from the scope of the invention. For example, the primary amino acid structures of the fusion proteins of the invention may be modified by creating amino acid mutants. Such mutants should retain more than 50% of the complement regulatory activity of the parent terminal CIP. Other modifications and variations include forming derivatives of the fusion protein to include covalent or aggregated conjugates of the protein or its fragments with other proteins or polypeptides. The following claims are intended to cover the specific embodiments set forth herein as well as such modifications, variations, and equivalents.

Throughout this application, various publications, patents, and patent applications have been referred to. The teachings and disclosures of these publications, patents, and patent applications in their entireties are hereby incorporated by reference into this application to more fully describe the state of the art to which the present invention pertains.

DEPOSITS

Plasmids pcDNA3/CD59-MCP-TM, pc8-hCD59-103, and pKS-/mCCPH, discussed above, have been deposited with the American Type Culture Collection, 12301 Parklawn Drive, Rockville, Maryland, 20852, United States of America, in

-43-

E. coli and have been assigned the designations 69530, 69231, and 69178, respectively. These deposits were made under the Budapest Treaty on the International Recognition of the Deposit of Micro-organisms for the Purposes of Patent Procedure (1977).

The deposits referred to above having ATCC accession numbers 69530, 69231, and 69178 were made on January 6, 1994, January 29, 1993, and January 6, 1993, respectively. Deposit 69530 was made in *Escherichia coli* strain TOP10F' which has the following genotype: F' {lacI^q TN10(Tet^R)} mcrA Δ(mrr-hsdRMS-mcrBC) φ80lacZΔM15 ΔlacX74 deoR recA1 araD139 Δ(ara-leu)7697 galU galK rpsL(Str^R) endA1 nupG. Deposits 69231 and 69178 were made in *Escherichia coli* strain DH5α which has the following genotype: F⁻ φ80dlacZΔM15 Δ(lacZYA-argF)U169 recA1 endA1 hsdR17(r_k⁻, m_k⁺) supE44 λ⁻ thi-1 gyrA96 relA1.

References

- 5 Ausubel, et al., eds., "Current Protocols in Molecular Biology", Wiley Interscience, John Wiley and Sons, New York, 1992.
- Albrecht, et al., 1992. Virology. 190:527-530.
- 10 Bhakdi, et al., 1991. Immunol. Today. 12: 318-321.
- Bradley. in Robertson. ed. "Teratocarcinomas and Embryonic Stem Cells a Practical Approach" IRL Press, Eynsham, Oxford, England, 1987.
- 15 Branden, et al., "Introduction to Protein Structure" Garland Publishing, Inc., New York, 1991.
- Brasile, et al., 1985. Transplantation. 40:672-675.
- 20 Brasile, et al., 1987. Trans. Proceed. 19:894-895.
- Brinster, et al., 1985. Proc. Natl. Acad. Sci. USA. 82:4438-4442.
- 25 Brown, et al., 1992. Cell. 68:533-544.
- Butikofer, et al., 1989. Blood. 74:1481-1485.
- Card, et al., 1991. J. Immunol. 146:4092-4098.
- 30 Cinek, et al., 1992. J. Immunol. 149:2262-2270.
- Cosset, et al., 1990. J. Virol. 64:1070-1078.
- 35 Dalmaso, et al., 1992. Am. J. Pathol. 140:1157-1166.
- Davies, et al., 1989. J. Exp. Med. 170:637-654.
- 40 Davitz, et al., 1987. Science. 238:81-84.
- Deckert, et al., 1992. J. Immunol. 148:672-677.
- Eglitis, et al., 1988. Biotechniques. 6:608-614.
- 45 Eikelenboom, et al., 1989. Virchow's Archiv. B. Cell. Pathol. 56:259-262.
- Engelman, et al., 1986. Annu. Rev. Biophys. Biophys. Chem. 15:321-353.
- 50 Esser. 1991. Immunol. Today. 12:316-318.

- Ferguson, et al., 1988. Ann. Rev. Biochem. 57:285-320.
- 5 Groux, et al., 1989. J. Immunol. 142:3013-3020.
- Gumley, et al., 1992. J. Immunol. 149:2615-2618.
- Harada, et al., 1990. J. Immunol. 144:1823-1828.
- 10 Hogan, et al., in "Manipulating the Mouse Embryo: A Laboratory Manual". Cold Spring Harbor Laboratory, Cold Spring Harbor, NY, 1986.
- Holguin, et al., 1989. J. Clin. Invest. 84:7-17.
- 15 Korman, et al., 1987, Proc. Natl. Acad. Sci. USA. 84:2150-2154.
- Kyte, et al., 1982. J. Mol. Biol. 157:105.
- 20 Lovell-Badge. in Robertson. ed. "Teratocarcinomas and Embryonic Stem Cells a Practical Approach" IRL Press, Eynsham, Oxford, England, 1987.
- 25 Lublin, et al., 1991. J. Exp. Med. 174:35-44.
- Markowitz, et al., 1988. J. Virol. 62:1120-1124.
- McMahon, et al., 1989. Cell. 58:1075-1084.
- 30 McMahon, et al., 1990 Cell. 62:1073-1085.
- Meri, et al., 1990. Immunology. 70:1-9.
- 35 Miller, et al., 1986. Mol. Cell Biol. 6:2895-2902.
- Miller, et al., 1989. Biotechniques. 7:981-990.
- Moran, et al., 1992. J. Immunol. 140:1736-1743.
- 40 Morgenstern, et al., 1990. Nucleic Acids Res. 18:3587-3596.
- Nakano, et al., 1993. Molec. Immunol. 30(suppl. 1):37.
- 45 Norris, et al., 1993. Blood. 82(Suppl.):202a.
- Nose, et al., 1990. Immunology. 70: 145-149.
- 50 Okada, et al., 1989a. Biochem. Biophys. Res. Commun. 162: 1553-1559.
- Okada, et al., 1989b. J. Immunol. 143:2262-2266.
- 55

- Pedersen, et al., 1990. "Transgenic Techniques in Mice - A Video Guide", Cold Spring Harbor Laboratory, Cold Spring Harbor, New York.
- 5 Petranka, et al., 1992. Proc. Natl. Acad. Sci. USA. 89:7876-7879.
- Petranka, et al., 1993. Molec. Immunol. 30(suppl. 1):44.
- 10 Philbrick, et al., 1990. Eur. J. Immunol. 20:87-92.
- Purcell, et al., 1991. Immunogenetics 33:335-344.
- 15 Robertson, et al., 1986. Nature. 323:445-448.
- Robertson. 1987. in Robertson. ed. "Teratocarcinomas and Embryonic Stem Cells a Practical Approach" IRL Press, Eynsham, Oxford, England.
- 20 Roldan, et al., 1990. EMBO J. 9:467-474.
- Rollins, et al., 1990. J. Immunol. 144:3478-3483.
- 25 Rollins, et al., 1991. J. Immunol. 146:2345-2351.
- Rother, et al., 1994. J. Virol. 68:730-737.
- 30 Sambrook, et al., Molecular cloning: a laboratory manual. Second Edition. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. 1989.
- Sandrin, et al., 1993. Proc. Natl. Acad. Sci. USA. 90:11391.
- 35 Sawada, et al., 1989. DNA Cell. Biol. 9:213-220.
- Schonermack, et al., 1986. J. Immunol. 136:1772.
- 40 Seaman, et al., 1991. J. Exp. Med. 173:251-260.
- Shevach, et al., 1989. Immunol. Today. 10:195-200.
- Sims, et al., 1989. J. Biol. Chem. 264:19228-19235.
- 45 Starzl, et al., 1993. Lancet. 341:65-71.
- Stefanova, et al., 1989. Mol. Immunol. 26:153-161.
- 50 Stefanova, et al., 1991. J. Immunol. 147:1587-1592.
- Su, et al., 1991. J. Cell Biol. 112:377-384.
- Sugita, et al., 1988. J. Biochem. 104:633-637.
- 55

- Suter, et al., 1992. J. Neurosci. 12:306-318.
- Talib, et al., 1991. Gene. 98:289-293.
- 5 Tone, et al., 1992. J. Mol. Biol. 227:971-976.
- Vakeva, et al., 1992. Lab. Invest. 67:608-616.
- 10 Venneker, et al., 1992. Exp. Clin. Immunogenet. 9:33- 47.
- Walsh, et al., 1991. Eur. J. Immunol. 21:847-850.
- 15 Warren, et al., 1987. Mol. Cell Biol. 7:1326-1332.
- Whitlow, et al., 1990. Cell Immunol. 126:176-184.
- Whitlow, et al., 1993. Blood. 81:510-516.
- 20 Williams, et al., 1988. Immunogenetics. 27:265-272.
- Wing, et al., 1992. Immunology. 76:140-145.
- 25 Yamada, et al., 1990. Neurosci. Lttrs. 112:161-166.
- Zalman, et al., 1986. Proc. Natl. Acad. Sci. USA. 83:6975-6979.
- 30 Zhao, et al., 1991. J. Biol. Chem. 266:13418-13422.

- 48 -

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: Rother, Russell
Rollins, Scott
Squinto, Stephen P
- (ii) TITLE OF INVENTION: Terminal Complement
Inhibitor Fusion Genes and Proteins
- (iii) NUMBER OF SEQUENCES: 14
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: Maurice M. Klee
 - (B) STREET: 1951 Burr Street
 - (C) CITY: Fairfield
 - (D) STATE: Connecticut
 - (E) COUNTRY: USA
 - (F) ZIP: 06430
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: 3.5 inch, 720 Kb storage
 - (B) COMPUTER: Dell 486/50
 - (C) OPERATING SYSTEM: DOS 6.2
 - (D) SOFTWARE: Word Perfect 6.0
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER:
 - (B) FILING DATE:
 - (C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER: 08/205,720
 - (B) FILING DATE: 3-MAR-1994
 - (C) CLASSIFICATION:

- 49 -

(viii) ATTORNEY/AGENT INFORMATION:

- (A) NAME: Klee, Maurice M.
- (B) REGISTRATION NUMBER: 30,399
- (C) REFERENCE/DOCKET NUMBER: ALX-129PCT

(ix) TELECOMMUNICATION INFORMATION:

- (A) TELEPHONE: (203) 255 1400
- (B) TELEFAX: (203) 254 1101

-50-

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 763 base pairs
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Double
- (D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: cDNA to mRNA

- (A) DESCRIPTION: BABCIP full length cDNA

(iii) HYPOTHETICAL: No

(iv) ANTI-SENSE: No

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Papio hamadryas

(vii) IMMEDIATE SOURCE:

- (A) LIBRARY: Baboon Spleen Lambda ZAPII cDNA
Library, Catalog # 936103,
Stratagene Cloning Systems,
La Jolla, California

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

```

GGTTATGTGC CCACACTTGC CTAGGCTGTG AATAGTTAGT ACCTCTGATT      50
ACTTAGTTAA ATATGCTTCT AGATGAGAAG TAGCGAAAGG CTGGAAGGGA      100
TCCCGGGCGC CGCCAGGTTC TGTGGACAAT CACA ATG GGA                140
                                   Met Gly
                                   -25

ATC CAA GGA GGG TCT GTC CTG TTC GGG CTG CTG CTT GTC CTG GCT 185
Ile Gln Gly Gly Ser Val Leu Phe Gly Leu Leu Leu Val Leu Ala
      -20                      -15                      -10

GTC TTC TGC CAT TCA GGT CAT AGC CTG CAG TGC TAC AAC TGT CCT 230
Val Phe Cys His Ser Gly His Ser Leu Gln Cys Tyr Asn Cys Pro
      -5                      1                      5

```

- 51 -

AAC	CCA	ACT	ACT	GAC	TGC	AAA	ACA	GCC	ATC	AAT	TGT	TCA	TCT	GGT	275
Asn	Pro	Thr	Thr	Asp	Cys	Lys	Thr	Ala	Ile	Asn	Cys	Ser	Ser	Gly	
		10					15					20			
TTT	GAT	ACG	TGT	CTC	ATT	GCC	AGA	GCT	GGG	TTA	CAA	GTA	TAT	AAC	320
Phe	Asp	Thr	Cys	Leu	Ile	Ala	Arg	Ala	Gly	Leu	Gln	Val	Tyr	Asn	
		25					30					35			
CAG	TGT	TGG	AAG	TTT	GCG	AAT	TGC	AAT	TTC	AAT	GAC	ATT	TCA	ACC	365
Gln	Cys	Trp	Lys	Phe	Ala	Asn	Cys	Asn	Phe	Asn	Asp	Ile	Ser	Thr	
		40					45					50			
CTC	TTG	AAG	GAA	AGC	GAG	CTA	CAG	TAC	TTC	TGC	TGC	AAG	AAG	GAC	410
Leu	Leu	Lys	Glu	Ser	Glu	Leu	Gln	Tyr	Phe	Cys	Cys	Lys	Lys	Asp	
		55					60					65			
CTG	TGT	AAC	TTT	AAC	GAA	CAG	CTT	GAA	AAT	GGT	GGG	ACA	TCC	TTA	455
Leu	Cys	Asn	Phe	Asn	Glu	Gln	Leu	Glu	Asn	Gly	Gly	Thr	Ser	Leu	
		70					75					80			
TCA	GAG	AAA	ACA	GTT	GTT	CTG	CTG	GTG	ACC	CTA	CTT	CTG	GCA	GCA	500
Ser	Glu	Lys	Thr	Val	Val	Leu	Leu	Val	Thr	Leu	Leu	Leu	Ala	Ala	
		85					90					95			
GCC	TGG	TGC	CTT	CAT	CCC	TAAGTCAACA	CCAGGAGAGC	TTCTCCCATA							548
Ala	Trp	Cys	Leu	His	Pro										
		100													
CTCCCCGTTT	CTGCGTAGTC	CCCTTTCCCT	CGTGCNGATT	CTAAAGGCTT											598
ATATTTTCCA	ACCGGATCCT	GTTGGGAAAG	AATAAAATTG	ACTTGAGCAA											648
CCTGGCTAAG	ATAGAGGGGC	TCTGGAAGAC	TTCGAAGACC	AGTCCTGTTT											698
GCAGGGAAGC	CCCACTTGAA	GGAAGAAGTT	TAAGAGTGAA	GTAGGTGTGA											748
CTTGAGCTAG	ATTGG														763

-52-

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 469 base pairs
 (B) TYPE: Nucleic Acid
 (C) STRANDEDNESS: Double
 (D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: cDNA to mRNA

- (A) DESCRIPTION: AGMCIP full length cDNA

(iii) HYPOTHETICAL: No

(iv) ANTI-SENSE: No

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Cercopithecus aethiops
 (H) CELL LINE: COS-1 (ATCC CRL 1650)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

```

                                TTCTGTGGAC AATCACA ATG GGA ATC   26
                                      Met Gly Ile
                                      -25

CAA GGA GGG TCT GTC CTG TTC GGG CTG CTG CTT GCC CTG GCT GTC   71
Gln Gly Gly Ser Val Leu Phe Gly Leu Leu Leu Ala Leu Ala Val
      -20                      -15                      -10

TTC TGC CAT TCA GGT CAT AGC CTG CAA TGC TAC AAC TGT CCT AAC 116
Phe Cys His Ser Gly His Ser Leu Gln Cys Tyr Asn Cys Pro Asn
      -5                      1                      5

CCA ACT ACT AAC TGC AAA ACA GCC ATC AAT TGT TCA TCT GGT TTT 161
Pro Thr Thr Asn Cys Lys Thr Ala Ile Asn Cys Ser Ser Gly Phe
      10                      15                      20

GAT ACG TGT CTC ATT GCC AGA GCT GGG TTA CAA GTA TAT AAC CAG 206
Asp Thr Cys Leu Ile Ala Arg Ala Gly Leu Gln Val Tyr Asn Gln
      25                      30                      35

TGT TGG AAG TTT GCG AAT TGC AAT TTC AAT GAC ATT TCA ACC CTC 251
Cys Trp Lys Phe Ala Asn Cys Asn Phe Asn Asp Ile Ser Thr Leu
      40                      45                      50

```

-53-

TTG	AAG	GAA	AGC	GAG	CTA	CAG	TAC	TTC	TGC	TGC	AAG	GAG	GAC	CTG	296
Leu	Lys	Glu	Ser	Glu	Leu	Gln	Tyr	Phe	Cys	Cys	Lys	Glu	Asp	Leu	
	55					60					65				
TGT	AAC	GAA	CAG	CTT	GAA	AAT	GGT	GGG	ACA	TCC	TTA	TCA	GAG	AAA	341
Cys	Asn	Glu	Gln	Leu	Glu	Asn	Gly	Gly	Thr	Ser	Leu	Ser	Glu	Lys	
	70					75					80				
ACA	GTT	CTT	CTG	CTG	GTG	ACC	CCA	CTT	CTG	GCA	GCA	GCC	TGG	TGC	386
Thr	Val	Leu	Leu	Leu	Val	Thr	Pro	Leu	Leu	Ala	Ala	Ala	Trp	Cys	
	85					90					95				
CTT	CAT	CCC	TAAGTCAACA	CCAGGAGAGC	TTCTCCCATATA	CTCCCCGTTC									435
Leu	His	Pro													
	100														
CTGCGTAGTC	CCCTTTCCCC	GGCCGCATTC	TAAA												469

- 54 -

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 396 base pairs
 (B) TYPE: Nucleic Acid
 (C) STRANDEDNESS: Double
 (D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: cDNA to mRNA

- (A) DESCRIPTION: SQMCIP full coding cDNA

(iii) HYPOTHETICAL: No

(iv) ANTI-SENSE: No

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Saimiri sciureus
 (H) CELL LINE: DPSO 114/74 (ATCC CCL 194)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

ATG	GGA	ATC	CAA	GGA	GGG	TCT	GTC	CTG	TTT	GGG	CTG	CTG	CTC	GTC	45
Met	Gly	Ile	Gln	Gly	Gly	Ser	Val	Leu	Phe	Gly	Leu	Leu	Leu	Val	
-25					-20					-15					
CTG	GCT	GTC	TTC	TGC	CAT	TCA	GGT	AAT	AGC	CTG	CAA	TGC	TAC	AGC	90
Leu	Ala	Val	Phe	Cys	His	Ser	Gly	Asn	Ser	Leu	Gln	Cys	Tyr	Ser	
-10					-5					1				5	
TGT	CCT	CTC	CCA	ACC	ATG	GAG	TCC	ATG	GAG	TGC	ACT	GCG	TCC	ACC	135
Cys	Pro	Leu	Pro	Thr	Met	Glu	Ser	Met	Glu	Cys	Thr	Ala	Ser	Thr	
				10					15					20	
AAC	TGT	ACA	TCT	AAT	CTT	GAT	TCG	TGT	CTC	ATT	GCC	AAA	GCC	GGG	180
Asn	Cys	Thr	Ser	Asn	Leu	Asp	Ser	Cys	Leu	Ile	Ala	Lys	Ala	Gly	
				25					30					35	
TCA	GGA	GTA	TAT	TAC	CGG	TGT	TGG	AAG	TTT	GAC	GAT	TGC	AGT	TTC	225
Ser	Gly	Val	Tyr	Tyr	Arg	Cys	Trp	Lys	Phe	Asp	Asp	Cys	Ser	Phe	
				40					45					50	
AAA	CGC	ATC	TCA	AAC	CAA	TTG	TCG	GAA	ACT	CAG	TTA	AAG	TAT	CAC	270
Lys	Arg	Ile	Ser	Asn	Gln	Leu	Ser	Glu	Thr	Gln	Leu	Lys	Tyr	His	
				55					60					65	

- 55 -

TGC	TGC	AAG	AAG	AAC	CTG	TGT	AAT	GTT	AAG	GAA	GTG	CTT	GAA	AAT	315
Cys	Cys	Lys	Lys	Asn	Leu	Cys	Asn	Val	Lys	Glu	Val	Leu	Glu	Asn	
				70					75					80	
GGT	GGG	ACA	ACC	TTA	TCA	AAG	AAA	ACA	ATT	CTT	CTG	CTG	GTG	ACC	360
Gly	Gly	Thr	Thr	Leu	Ser	Lys	Lys	Thr	Ile	Leu	Leu	Leu	Val	Thr	
				85					90					95	
CCG	TTT	CTG	GCA	GCA	GCC	TGG	AGC	CGT	CAT	CCC	TAA				396
Pro	Phe	Leu	Ala	Ala	Ala	Trp	Ser	Arg	His	Pro					
				100											

-56-

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 387 base pairs
 (B) TYPE: Nucleic Acid
 (C) STRANDEDNESS: Double
 (D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: cDNA to mRNA

- (A) DESCRIPTION: OWMCIP full coding cDNA

(iii) HYPOTHETICAL: No

(iv) ANTI-SENSE: No

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Aotus trivirgatus
 (H) CELL LINE: OMK (ATCC CRL 1556)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

ATG GGA ATT CAA GGA GGG TCT GTC CTG TTT GGG CTG CTG CTC GTC	45
Met Gly Ile Gln Gly Gly Ser Val Leu Phe Gly Leu Leu Leu Val	
-25 -20 -15	
CTG GCT GTC TTC TGC CAT TCA GGT AAT AGC CTG CAG TGC TAC AGC	90
Leu Ala Val Phe Cys His Ser Gly Asn Ser Leu Gln Cys Tyr Ser	
-10 -5 1 5	
TGT CCT TAC CCA ACC ACT CAG TGC ACT ATG ACC ACC AAC TGT ACA	135
Cys Pro Tyr Pro Thr Thr Gln Cys Thr Met Thr Thr Asn Cys Thr	
10 15 20	
TCT AAT CTT GAT TCG TGT CTC ATT GCC AAA GCC GGG TCA CGA GTA	180
Ser Asn Leu Asp Ser Cys Leu Ile Ala Lys Ala Gly Ser Arg Val	
25 30 35	
TAT TAC CGG TGT TGG AAG TTT GAG GAT TGC ACT TTC AGC CGC GTT	225
Tyr Tyr Arg Cys Trp Lys Phe Glu Asp Cys Thr Phe Ser Arg Val	
40 45 50	
TCA AAC CAA TTG TCT GAA AAT GAG TTA AAG TAT TAC TGC TGC AAG	270
Ser Asn Gln Leu Ser Glu Asn Glu Leu Lys Tyr Tyr Cys Cys Lys	
55 60 65	

-57-

AAG	AAC	CTG	TGT	AAC	TTT	AAT	GAA	GCG	CTT	AAA	AAT	GGT	GGG	ACA	315
Lys	Asn	Leu	Cys	Asn	Phe	Asn	Glu	Ala	Leu	Lys	Asn	Gly	Gly	Thr	
				70					75					80	
ACC	TTA	TCA	AAG	AAA	ACA	GTC	CTC	CTG	CTG	GTG	ATC	CCA	TTT	CTG	360
Thr	Leu	Ser	Lys	Lys	Thr	Val	Leu	Leu	Leu	Val	Ile	Pro	Phe	Leu	
				85					90					95	
GTA	GCA	GCC	TGG	AGC	CTT	CAT	CCC	TAA							387
Val	Ala	Ala	Trp	Ser	Leu	His	Pro								
				100											

-58-

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 387 base pairs
 (B) TYPE: Nucleic Acid
 (C) STRANDEDNESS: Double
 (D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: cDNA to mRNA

- (A) DESCRIPTION: MARCIP full coding cDNA

(iii) HYPOTHETICAL: No

(iv) ANTI-SENSE: No

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: *Saguinus nigricollis*
 (H) CELL LINE: 1283.Lu (ATCC CRL 6297)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

ATG	GGA	ATC	CAA	GGA	GGG	TCT	GTC	CTG	TTT	GGG	CTG	CTG	CTC	ATC	45
Met	Gly	Ile	Gln	Gly	Gly	Ser	Val	Leu	Phe	Gly	Leu	Leu	Leu	Ile	
-25					-20					-15					
CTG	GCT	GTC	TTC	TGC	CAT	TCA	GGT	CAT	AGC	CTG	CAG	TGC	TAC	AGC	90
Leu	Ala	Val	Phe	Cys	His	Ser	Gly	His	Ser	Leu	Gln	Cys	Tyr	Ser	
-10					-5					1				5	
TGT	CCT	TAC	TCA	ACC	GCT	CGG	TGC	ACT	ACG	ACC	ACC	AAC	TGT	ACA	135
Cys	Pro	Tyr	Ser	Thr	Ala	Arg	Cys	Thr	Thr	Thr	Thr	Asn	Cys	Thr	
				10					15					20	
TCT	AAT	CTT	GAT	TCA	TGT	CTC	ATT	GCC	AAA	GCC	GGG	TTA	CGA	GTA	180
Ser	Asn	Leu	Asp	Ser	Cys	Leu	Ile	Ala	Lys	Ala	Gly	Leu	Arg	Val	
				25					30					35	
TAT	TAC	CGG	TGT	TGG	AAG	TTT	GAG	GAT	TGC	ACT	TTC	AGA	CAA	CTT	225
Tyr	Tyr	Arg	Cys	Trp	Lys	Phe	Glu	Asp	Cys	Thr	Phe	Arg	Gln	Leu	
				40					45					50	
TCA	AAC	CAA	TTG	TCG	GAA	AAT	GAG	TTA	AAG	TAT	CAC	TGC	TGC	AGG	270
Ser	Asn	Gln	Leu	Ser	Glu	Asn	Glu	Leu	Lys	Tyr	His	Cys	Cys	Arg	
				55					60					65	

-59-

GAG	AAC	CTG	TGT	AAC	TTT	AAC	GGA	ATA	CTT	GAA	AAT	GGT	GGG	ACA	315
Glu	Asn	Leu	Cys	Asn	Phe	Asn	Gly	Ile	Leu	Glu	Asn	Gly	Gly	Thr	
				70					75					80	
ACC	TTA	TCA	AAG	AAA	ACA	GTT	CTT	CTG	CTG	GTG	ACC	CCT	TTT	CTG	360
Thr	Leu	Ser	Lys	Lys	Thr	Val	Leu	Leu	Leu	Val	Thr	Pro	Phe	Leu	
				85					90					95	
GCA	GCA	GCC	TGG	AGC	CTT	CAT	CCC	TAA							387
Ala	Ala	Ala	Trp	Ser	Leu	His	Pro								
				100											

-60-

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1039 base pairs
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Double
- (D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: cDNA to mRNA

- (A) DESCRIPTION: HVS-15 full length cDNA

(iii) HYPOTHETICAL: No

(iv) ANTI-SENSE: No

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Herpesvirus saimiri

(x) PUBLICATION INFORMATION:

- (A) AUTHORS: Albrecht, J.C.
Nicholas, J.
Cameron, K.R.
Newman, C.
Fleckenstein, B.
Honess, R.W.

- (B) TITLE: Herpesvirus samiri has a gene specifying
a homologue of the cellular membrane
glycoprotein CD59.

- (C) JOURNAL: Virology
- (D) VOLUME: 190
- (F) PAGES: 527-530
- (G) DATE: 1992

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

```
AAGCTTCTAT TTATACTACA TTAGAGGCAT TTTTTCAAAA GCAAAAATGC      50
CTCTAATTAT ATACACTGTA CTATTACCT CTATTACACA TTTTCTATTT      100
TAAGTCTGAT AGTGATTAAT CAAGAAAAAA GTTTGTGGTT CTCAGGGGAT      150
TAGTTCACAA GCTGTCTGAG GTTAAGGGTG TTTCTTTGGC ACTGACACAG      200
AAGTTGCTAT AAGAATTGAA GCTTGCTTTA CAAAAAGTTA CTTGTGATTA      250
```

[illegible]

- 62 -

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1139 base pairs
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Double
- (D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: cDNA to mRNA

- (A) DESCRIPTION: CD59 full length cDNA

(iii) HYPOTHETICAL: No

(iv) ANTI-SENSE: No

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(x) PUBLICATION INFORMATION:

- (A) AUTHORS: Philbrick, W.M.
Palfree, R.G.E.
Maher, S.E.
Bridgett, M.M.
Sirlin S.
Bothwell, A.L.M.
- (B) TITLE: The CD59 antigen is a structural
homologue of murine Ly-6 antigens but
lacks interferon inducibility.
- (C) JOURNAL: European Journal of Immunology
- (D) VOLUME: 20
- (F) PAGES: 87-92
- (G) DATE: JAN-1990

- 63 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

CGCAGAAGCG GCTCGAGGCT GGAAGAGGAT CCTGGGCGCC GCAGGTTCTG	50
TGGACAATCA CA ATG GGA ATC CAA GGA GGG TCT GTC CTG TTC	92
Met Gly Ile Gln Gly Gly Ser Val Leu Phe	
-25 -20	
GGG CTG CTG CTC GTC CTG GCT GTC TTC TGC CAT TCA GGT CAT	134
Gly Leu Leu Leu Val Leu Ala Val Phe Cys His Ser Gly His	
-15 -10 -5	
AGC CTG CAG TGC TAC AAC TGT CCT AAC CCA ACT GCT GAC TGC	176
Ser Leu Gln Cys Tyr Asn Cys Pro Asn Pro Thr Ala Asp Cys	
1 5 10	
AAA ACA GCC GTC AAT TGT TCA TCT GAT TTT GAT GCG TGT CTC	218
Lys Thr Ala Val Asn Cys Ser Ser Asp Phe Asp Ala Cys Leu	
15 20 25	
ATT ACC AAA GCT GGG TTA CAA GTG TAT AAC AAG TGT TGG AAG	260
Ile Thr Lys Ala Gly Leu Gln Val Tyr Asn Lys Cys Trp Lys	
30 35 40	
TTT GAG CAT TGC AAT TTC AAC GAC GTC ACA ACC CGC TTG AGG	302
Phe Glu His Cys Asn Phe Asn Asp Val Thr Thr Arg Leu Arg	
45 50 55	
GAA AAT GAG CTA ACG TAC TAC TGC TGC AAG AAG GAC CTG TGT	344
Glu Asn Glu Leu Thr Tyr Tyr Cys Cys Lys Lys Asp Leu Cys	
60 65	
AAC TTT AAC GAA CAG CTT GAA AAT GGT GGG ACA TCC TTA TCA	386
Asn Phe Asn Glu Gln Leu Glu Asn Gly Gly Thr Ser Leu Ser	
70 75 80	
GAG AAA ACA GTT CTT CTG CTG GTG ACT CCA TTT CTG GCA GCA	428
Glu Lys Thr Val Leu Leu Leu Val Thr Pro Phe Leu Ala Ala	
85 90 95	
GCC TGG AGC CTT CAT CCC TAA G TCAACACCAG GAGAGCTTCT	470
Ala Trp Ser Leu His Pro	
100	
CCCAAACCTCC CCGTTCCTGC GTAGTCCGCT TTCTCTTGCT GCCACATTCT	520
AAAGGCTTGA TATTTTCCAA ATGGATCCTG TTGGGAAAGA ATAAAATTAG	570
CTTGAGCAAC CTGGCTAAGA TAGAGGGGTC TGGGAGACTT TGAAGACCAG	620
TCCTGCCCCGC AGGGAAGCCC CACTTGAAGG AAGAAGTCTA AGAGTGAAGT	670
AGGTGTGACT TGAAC TAGAT TGCATGCTTC CTCCTTTGCT CTTGGGAAGA	720

- 64 -

CCAGCTTTGC	AGTGACAGCT	TGAGTGGGTT	CTCTGCAGCC	CTCAGATTAT	770
TTTTCCTCTG	GCTCCTTGGA	TGTAGTCAGT	TAGCATCATT	AGTACATCTT	820
TGGAGGGTGG	GGCAGGAGTA	TATGAGCATC	CTCTCTCACA	TGGAACGCTT	870
TCATAAACTT	CAGGGATCCC	GTGTTGCCAT	GGAGGCATGC	CAAATGTTCC	920
ATATGTGGGT	GTCAGTCAGG	GACAACAAGA	TCCTTAATGC	AGAGCTAGAG	970
GACTTCTGGC	AGGGAAGTGG	GGAAGTGTTT	CAGATTCCAG	ATAGCAGGGC	1020
ATGAAAACCTT	AGAGAGGTAC	AAGTGGCTGA	AAATCGAGTT	TTTCCTCTGT	1070
CTTTAAATTT	TATATGGGCT	TTGTTATCTT	CCACTGGAAA	AGTGTAATAG	1120
CATACATCAA	TGGTGTGTT				1139

-65-

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1530 base pairs
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Double
- (D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: cDNA to mRNA

- (A) DESCRIPTION: MCP (CD46) full length cDNA

(iii) HYPOTHETICAL: No

(iv) ANTI-SENSE: No

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(x) PUBLICATION INFORMATION:

- (A) AUTHORS: Lublin, D.M.
Liszewski, M.K.
Post, T.W.
Arce, M.A.
LeBeau, M.M.
Rebentisch, M.B.
Lemons, R.S.
Seya, T.
Atkinson, J.P.
- (B) TITLE: Molecular cloning and Chromosomal
Localization of Membrane Cofactor
Protein (MCP): Evidence for
Inclusion in the Multi-Gene Family
of Complement-Regulatory Proteins.
- (C) JOURNAL: Journal of Experimental Medicine
- (D) VOLUME: 168
- (F) PAGES: 181-194
- (G) DATE: 1988

- 66 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

TCTGCTTTCC	TCCGGAGAAA	TAACAGCGTC	TTCCGCGCCG	CGC	ATG	GAG	49
					Met	Glu	
CCT CCC GGC CGC CGC GAG TGT CCC TTT CCT TCC TGG CGC TTT	91						
Pro Pro Gly Arg Arg Glu Cys Pro Phe Pro Ser Trp Arg Phe							
-30 -25 -20							
CCT GGG TTG CTT CTG GCG GCC ATG GTG TTG CTG CTG TAC TCC	133						
Pro Gly Leu Leu Leu Ala Ala Met Val Leu Leu Leu Tyr Ser							
-15 -10 -5							
TTC TCC GAT GCC TGT GAG GAG CCA CCA ACA TTT GAA GCT ATG	175						
Phe Ser Asp Ala Cys Glu Glu Pro Pro Thr Phe Glu Ala Met							
1 5 10							
GAG CTC ATT GGT AAA CCA AAA CCC TAC TAT GAG ATT GGT GAA	217						
Glu Leu Ile Gly Lys Pro Lys Pro Tyr Tyr Glu Ile Gly Glu							
15 20							
CGA GTA GAT TAT AAG TGT AAA AAA GGA TAC TTC TAT ATA CCT	259						
Arg Val Asp Tyr Lys Cys Lys Lys Gly Tyr Phe Tyr Ile Pro							
25 30 35							
CCT CTT GCC ACC CAT ACT ATT TGT GAT CGG AAT CAT ACA TGG	301						
Pro Leu Ala Thr His Thr Ile Cys Asp Arg Asn His Thr Trp							
40 45 50							
CTA CCT GTC TCA GAT GAC GCC TGT TAT AGA GAA ACA TGT CCA	343						
Leu Pro Val Ser Asp Asp Ala Cys Tyr Arg Glu Thr Cys Pro							
55 60 65							
TAT ATA CGG GAT CCT TTA AAT GGC CAA GCA GTC CCT GCA AAT	385						
Tyr Ile Arg Asp Pro Leu Asn Gly Gln Ala Val Pro Ala Asn							
70 75 80							
GGG ACT TAC GAG TTT GGT TAT CAG ATG CAC TTT ATT TGT AAT	427						
Gly Thr Tyr Glu Phe Gly Tyr Gln Met His Phe Ile Cys Asn							
85 90							
GAG GGT TAT TAC TTA ATT GGT GAA GAA ATT CTA TAT TGT GAA	469						
Glu Gly Tyr Tyr Leu Ile Gly Glu Glu Ile Leu Tyr Cys Glu							
95 100 105							
CTT AAA GGA TCA GTA GCA ATT TGG AGC GGT AAG CCC CCA ATA	511						
Leu Lys Gly Ser Val Ala Ile Trp Ser Gly Lys Pro Pro Ile							
110 115 120							
TGT GAA AAG GTT TTG TGT ACA CCA CCT CCA AAA ATA AAA AAT	553						
Cys Glu Lys Val Leu Cys Thr Pro Pro Pro Lys Ile Lys Asn							
125 130 135							

- 67 -

GGA	AAA	CAC	ACC	TTT	AGT	GAA	GTA	GAA	GTA	TTT	GAG	TAT	CTT	595
Gly	Lys	His	Thr	Phe	Ser	Glu	Val	Glu	Val	Phe	Glu	Tyr	Leu	
			140					145					150	
GAT	GCA	GTA	ACT	TAT	AGT	TGT	GAT	CCT	GCA	CCT	GGA	CCA	GAT	637
Asp	Ala	Val	Thr	Tyr	Ser	Cys	Asp	Pro	Ala	Pro	Gly	Pro	Asp	
				155					160					
CCA	TTT	TCA	CTT	ATT	GGA	GAG	AGC	ACG	ATT	TAT	TGT	GGT	GAC	679
Pro	Phe	Ser	Leu	Ile	Gly	Glu	Ser	Thr	Ile	Tyr	Cys	Gly	Asp	
165					170					175				
AAT	TCA	GTG	TGG	AGT	CGT	GCT	GCT	CCA	GAG	TGT	AAA	GTG	GTC	721
Asn	Ser	Val	Trp	Ser	Arg	Ala	Ala	Pro	Glu	Cys	Lys	Val	Val	
	180					185					190			
AAA	TGT	CGA	TTT	CCA	GTA	GTC	GAA	AAT	GGA	AAA	CAG	ATA	TCA	763
Lys	Cys	Arg	Phe	Pro	Val	Val	Glu	Asn	Gly	Lys	Gln	Ile	Ser	
		195					200					205		
GGA	TTT	GGA	AAA	AAA	TTT	TAC	TAC	AAA	GCA	ACA	GTT	ATG	TTT	805
Gly	Phe	Gly	Lys	Lys	Phe	Tyr	Tyr	Lys	Ala	Thr	Val	Met	Phe	
			210					215					220	
GAA	TGC	GAT	AAG	GGT	TTT	TAC	CTC	GAT	GGC	AGC	GAC	ACA	ATT	847
Glu	Cys	Asp	Lys	Gly	Phe	Tyr	Leu	Asp	Gly	Ser	Asp	Thr	Ile	
				225					230					
GTC	TGT	GAC	AGT	AAC	AGT	ACT	TGG	GAT	CCC	CCA	GTT	CCA	AAG	889
Val	Cys	Asp	Ser	Asn	Ser	Thr	Trp	Asp	Pro	Pro	Val	Pro	Lys	
235					240					245				
TGT	CTT	AAA	GTG	TCG	ACT	TCT	TCC	ACT	ACA	AAA	TCT	CCA	GCG	931
Cys	Leu	Lys	Val	Ser	Thr	Ser	Ser	Thr	Thr	Lys	Ser	Pro	Ala	
	250					255					260			
TCC	AGT	GCC	TCA	GGT	CCT	AGG	CCT	ACT	TAC	AAG	CCT	CCA	GTC	973
Ser	Ser	Ala	Ser	Gly	Pro	Arg	Pro	Thr	Tyr	Lys	Pro	Pro	Val	
		265					270					275		
TCA	AAT	TAT	CCA	GGA	TAT	CCT	AAA	CCT	GAG	GAA	GGA	ATA	CTT	1015
Ser	Asn	Tyr	Pro	Gly	Tyr	Pro	Lys	Pro	Glu	Glu	Gly	Ile	Leu	
			280				285						290	
GAC	AGT	TTG	GAT	GTT	TGG	GTC	ATT	GCT	GTG	ATT	GTT	ATT	GCC	1057
Asp	Ser	Leu	Asp	Val	Trp	Val	Ile	Ala	Val	Ile	Val	Ile	Ala	
				295					300					
ATA	GTT	GTT	GGA	GTT	GCA	GTA	ATT	TGT	GTT	GTC	CCG	TAC	AGA	1099
Ile	Val	Val	Gly	Val	Ala	Val	Ile	Cys	Val	Val	Pro	Tyr	Arg	
305					310					315				
TAT	CTT	CAA	AGG	AGG	AAG	AAG	AAA	GGG	AAA	GCA	GAT	GGT	GGA	1141
Tyr	Leu	Gln	Arg	Arg	Lys	Lys	Lys	Gly	Lys	Ala	Asp	Gly	Gly	
	320					325					330			

- 68 -

GCT GAA TAT GCC ACT TAC CAG ACT AAA TCA ACC ACT CCA GCA 1183
Ala Glu Tyr Ala Thr Tyr Gln Thr Lys Ser Thr Thr Pro Ala
335 340 345

GAG CAG AGA GGC TGA AT AGATTCCACA ACCTGGTTTG CCAGTTCATC 1230
Glu Gln Arg Gly
350

TTTTGACTCT ATTAAAATCT TCAATAGTTG TTATTCTGTA GTTTCACTCT 1280

CATGAGTGCA ACTGTGGCTT AGCTAATATT GCAATGTGGC TTGAATGTAG 1330

GTAGCATCCT TTGATGCTTC TTTGAAACTT GTATGAATTT GGGTATGAAC 1380

AGATTGCCTG CTTTCCCTTA AATAACACTT AGATTTATTG GACCAGTCAG 1430

CACAGCATGC CTGGTTGTAT TAAAGCAGGG ATATGCTGTA TTTTATAAAA 1480

TTGGCAAAAT TAGAGAAATA TAGTTCACAA TGAAATTATA TTTTCTTTGT 1530

-69-

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 bases
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

- (ii) MOLECULE TYPE: Other nucleic acid
- (iii) HYPOTHETICAL: No
- (iv) ANTI-SENSE: No

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

CGCGAGGCCT ACTTACAAGC CTCCAG

26

(2) INFORMATION FOR SEQ ID NO 10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 bases
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

- (ii) MOLECULE TYPE: Other nucleic acid
- (iii) HYPOTHETICAL: No
- (iv) ANTI-SENSE: Yes

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

CGCGCTATTC AGCCTCTCTG CTCTGC

26

- 70 -

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 261 base pairs
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Double
- (D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Other nucleic acid

- (A) DESCRIPTION: MCP PCR Product

(iii) HYPOTHETICAL: No

(iv) ANTI-SENSE: No

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

```
CGCGAGGCCT ACTTACAAGC CTCCAGTCTC AAATTATCCA GGATATCCTA   50
AACCTGAGGA AGGAATACTT GACAGTTTGG ATGTTTGGGT CATTGCTGTG  100
ATTGTTATTG CCATAGTTGT TGGAGTTGCA GTAATTGTG TTGTCCCGTA  150
CAGATATCTT CAAAGGAGGA AGAAGAAAGG GAAAGCAGAT GGTGGAGCTG  200
AATATGCCAC TTACCAGACT AAATCAACCA CTCCAGCAGA GCAGAGAGGC  250
TGAATAGCGC G                                           261
```


-71-

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 69 bases
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Other nucleic acid

(iii) HYPOTHETICAL: No

(iv) ANTI-SENSE: No

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

CCGGACCTGT GTAAC TTAA CGAACAGCTT GAAAATATTG GTAGGATATG 50
CAATGGAAAT TGT TACAAC 69

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 bases
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Other nucleic acid

(iii) HYPOTHETICAL: No

(iv) ANTI-SENSE: Yes

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

TAGTTACTGC CCGGACATGC 20

- 72 -

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 264 base pairs
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Double
- (D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Other nucleic acid

(A) DESCRIPTION: CCPH PCR Product

(iii) HYPOTHETICAL: No

(iv) ANTI-SENSE: No

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

```
CCGGACCTGT GTAAC TT TAA CGAACAGCTT GAAAATATTG GTAGGATATG      50
CAATGGAAAT TGTACA ACTA GCATGCCCCAC TCAAACATAT ACAATAATTA      100
CTGCGCGCTA TACAAGTCAC ATATATTTCC CTACTGGGAA AACCTATAAA      150
CTTCCTCGGG GAGTTCTAGT AATTATTCTT ACCACAAGCT TTATTATTAT      200
TGAATAATA CTTACTGGAG TGTGTTTACA TAGGTGCAGA GTGTGCATGT      250
CCGGGCAGTA ACTA                                         264
```

What is claimed is:

1. A nucleic acid molecule comprising:
 - (a) a sequence encoding a chimeric protein which comprises:
 - (i) a first polypeptide region comprising a portion of a parent terminal complement inhibitor protein, said portion including a complete Ly-6 motif and not including an operative signal sequence directing the attachment of a glycosyl-phosphatidylinositol (GPI) anchor; and
 - (ii) a second polypeptide region linked to the first polypeptide region, said second polypeptide region comprising a transmembrane domain from a heterologous protein; or
 - (b) a sequence complementary to (a); or
 - (c) both (a) and (b)said molecule being substantially free of nucleic acid molecules not containing (a), (b), or (c).
2. The nucleic acid molecule of Claim 1 wherein said chimeric protein has greater than 50% of the complement inhibitory activity of the parent terminal complement inhibitor protein.
3. The nucleic acid molecule of Claim 1 wherein the portion of the parent terminal complement inhibitor protein comprises said protein minus amino acid residues downstream of its Ly-6 motif.
4. The nucleic acid molecule of Claim 1 wherein the chimeric protein has complement inhibitory activity against human complement.
5. A nucleic acid vector comprising the nucleic acid molecule of Claim 1 operatively linked to a second nucleic acid molecule so that a host containing the vector expresses the chimeric protein.
6. A recombinant host containing the vector of Claim 4.

- 74 -

7. A process for protecting a non-human organ from human complement attack comprising introducing the nucleic acid molecule of Claim 4 into a pluripotent cell capable of producing a non-human transgenic animal and producing the non-human transgenic animal from said cell, whereby the resistance of an organ of said non-human transgenic animal to human complement attack is enhanced.

8. Cells isolated from the transgenic animal of Claim 7.

9. A chimeric protein comprising:

(i) a first polypeptide region comprising a portion of a parent terminal complement inhibitor protein, said portion including a complete Ly-6 motif and not including an operative signal sequence directing the attachment of a glycosyl-phosphatidylinositol (GPI) anchor; and

(ii) a second polypeptide region linked to the first polypeptide region, said second polypeptide region comprising a transmembrane domain from a heterologous protein.

10. The chimeric protein of Claim 9 wherein said protein has greater than 50% of the complement inhibitory activity of the parent terminal complement inhibitor protein.

11. The chimeric protein of Claim 9 wherein the portion of the parent terminal complement inhibitor protein comprises said parent protein minus amino acid residues downstream of its Ly-6 motif.

12. The chimeric protein of Claim 9 wherein the protein has complement inhibitory activity against human complement.

13. A membrane bound terminal complement inhibitor protein which is substantially resistant to the action of lipase enzymes which can cleave a glycosyl-phosphatidylinositol (GPI) anchor.

14. The membrane bound terminal complement inhibitor protein of Claim 13 wherein the protein comprises a Ly-6 motif.

1/6

CD59	MG1	QGG	SVL	FGL	LLV	LAV	FCH	SGH	S	LQ	CYN	CPN	PTA	---	DCK	
AGMCIP	MG1	QGG	SVL	FGL	LLA	LAV	FCH	SGH	S	LQ	CYN	CPN	PTT	---	NCK	
BABCIP	MG1	QGG	SVL	FGL	LLV	LAV	FCH	SGH	S	LQ	CYN	CPN	PTT	---	DCK	
OWMCIP	MG1	QGG	SVL	FGL	LLV	LAV	FCH	SGN	S	LQ	CYS	CPY	PTT	---	QCT	
MARCIP	MG1	QGG	SVL	FGL	LLI	LAV	FCH	SGH	S	LQ	CYS	CPY	STA	---	RCT	
SQMCIP	MG1	QGG	SVL	FGL	LLV	LAV	FCH	SGN	S	LQ	CYS	CPL	PTM	ESM	ECT	
HVS-15				M	YIL	FTL	VLT	F-V	FCK	PIH	S	LQ	CSH	STM	---	QCT

CD59	TAV	NCS	SDF	DAC	LIT	KAG	LQV	YNK	CWK	FEH	CNF	NDV	TTR	LRE	NEL
AGMCIP	TAI	NCS	SGF	DTC	LIA	RAG	LQV	YNQ	CWK	FAN	CNF	NDI	STL	LKE	SEL
BABCIP	TAI	NCS	SGF	DTC	LIA	RAG	LQV	YNQ	CWK	FAN	CNF	NDI	STL	LKE	SEL
OWMCIP	MTT	NCT	SNL	DSC	LIA	KAG	SRV	YYR	CWK	FED	CTF	SRV	SNQ	LSE	NEL
MARCIP	TTT	NCT	SNL	DSC	LIA	KAG	LRV	YYR	CWK	FED	CTF	RQL	SNQ	LSE	NEL
SQMCIP	AST	NCT	SNL	DSC	LIA	KAG	SGV	YYR	CWK	FDD	CSF	KRI	SNQ	LSE	TQL
HVS-15	TST	SCT	SNL	DSC	LIA	KAG	SGV	YYR	CWK	FDD	CSF	KRI	SNQ	LSE	TQL

CD59	TYV	CCK	KDL	CNF	NEQ	LEN	GGT	SLS	EKT	VLL	LVT	PFL	AAA	WSL	HP
AGMCIP	QYF	CCK	EDL	CN-	-EQ	LEN	GGT	SLS	EKT	VLL	LVT	PLL	AAA	WCL	HP
BABCIP	QYF	CCK	KDL	CNF	NEQ	LEN	GGT	SLS	EKT	VVL	LVT	LLL	AAA	WCL	HP
OWMCIP	KYV	CCK	KNL	CNF	NEA	LKN	GGT	TLS	KKT	VLL	LVI	PFL	VAA	WSL	HP
MARCIP	KYH	CCK	ENL	CNF	NGI	LEN	GGT	TLS	KKT	VLL	LVT	PFL	AAA	WSL	HP
SQMCIP	KYH	CCK	KNL	CNV	KEV	LEN	GGT	TLS	KKT	ILL	LVT	PFL	AAA	WSR	HP
HVS-15	KYH	CCK	KNL	CNV	NKG	IEN	IKR	TIS	DKA	LLL	LLA	LFL	VTA	WNF	PL

Fig. 1

2/6




Fig. 2

3/6




Fig. 3

4/6




Fig. 4A




Fig. 4B

5/6




Fig. 5

6/6




Fig. 6

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US95/02944

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : Please See Extra Sheet.

US CL : 424/93.21; 435/41, 69.1, 69.7, 172.1, 172.3, 240.1, 240.21, 243; 514/44; 530/380; 536/23.4

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/93.21; 435/41, 69.1, 69.7, 172.1, 172.3, 240.1, 240.21, 243; 514/44; 530/380; 536/23.4

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	HUMAN GENE THERAPY, Volume 2, issued 1991, F.D. Ledley, "Clinical considerations in the design of protocols for somatic gene therapy", pages 77-83, see entire document.	6-8
Y	THE LANCET, Volume 339, issued 21 March 1992, A.A. Gutierrez et al., "Gene therapy for Cancer", pages 715-721, see entire article.	6-8
X,P	BLOOD, Volume 84, Number 8, issued 15 October 1994, R.P. Rother et al., "Expression of recombinant transmembrane CD59 in paroxysmal nocturnal hemoglobinuria B cells confers resistance to human complement", pages 2604-2611, see entire document.	1-14

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z*	document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means		
P document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search 30 MAY 1995	Date of mailing of the international search report 07 JUN 1995
--	---

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

BRIAN R. STANTON

Telephone No. (703) 308-0196

Form PCT/ISA/210 (second sheet)(July 1992)*

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US95/02944

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EXPERIMENTAL AND CLINICAL IMMUNOGENETICS, Volume 9, issued 1992, G.T. Venneker et al., "CD59: A molecule involved in antigen presentation as well as downregulation of membrane attack complex", pages 33-47, see entire document.	1-14
Y	EUROPEAN JOURNAL OF IMMUNOLOGY, Volume 21, issued 1991, L.A. Walsh et al., "Transfection of human CD59 complementary DNA into rat cells confers resistance to human complement", pages 847-850, see entire document.	1-14
Y	CELL, Volume 73, issued 21 May 1993, Takeda et al., "Deficiency of the GPI anchor caused by a somatic mutation of the PIG-A gene in paroxysmal nocturnal hemoglobinuria", pages 703-711, see entire document.	1-14
Y	JOURNAL OF EXPERIMENTAL MEDICINE, Volume 174, issued July 1991, D.M. Lublin et al., "Phospholipid-anchored and transmembrane versions of either decay-accelerating factor or membrane cofactor protein show equal efficiency in protection from complement-mediated cell damage", pages 35-44, see entire document.	1-14
Y	DNA AND CELL BIOLOGY, Volume 9, Number 3, issued 1990, R. Sawada et al., "Isolation and expression of the full-length cDNA encoding CD59 antigen of human lymphocytes", pages 213-220, see entire document.	1-14
Y	SEMINARS IN HEMATOLOGY, Volume 26, Number 3, issued July 1989, B. Rotoli et al., "Paroxysmal nocturnal hemoglobinuria", pages 201-207, see entire document.	1-14

Form PCT/ISA/210 (continuation of second sheet)(July 1992)*

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US95/02944

A. CLASSIFICATION OF SUBJECT MATTER:

IPC (6):

AO1N 63/00; A61K 35/14, 38/00; C07H 17/00; C07K 14/00; C12N 1/00, 5/00, 5/06, 5/22, 7/01, 15/00, 15/03, 15/09, 15/06, 15/11; C12P 1/00, 21/06

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

Databases: APS, Medline, CA, CAPreviews, Embase, Derwent, Biosis

Search Terms: chimera?; protein?; terminal; complement; inhibitor?; ly-6; ly6; glycosyl?; phosphatidylinositol; gpi; transmembran?; anchor; motif; cysteine?; vector?; lysis; miri; cd59; daf; transgen?; lipase?; rother?/au; rollins?/au; squinto?/au

THIS PAGE BLANK (USPTO)